

# EURO4M: European Reanalysis and Observations for Monitoring

Proposal to the Seventh Framework Programme  
Cooperation Theme 9 “SPACE”

Proposal full title:	European Reanalysis and Observations for Monitoring
Proposal acronym:	EURO <sub>4</sub> M
Type of funding scheme:	Collaborative project (small-medium scale focused research project)
Work programme topics addressed:	SPA.2009.1.1.02 Monitoring of climate change issues
Name of coordinating person:	Dr. Albert M.G. Klein Tank (KNMI)
List of participants:	See below

No.	Participant organisation name	Short	Country
1	Royal Netherlands Meteorological Institute	KNMI	Netherlands
2	Met Office	MO	United Kingdom
3	University Rovira i Virgili	URV	Spain
4	National Meteorological Administration	NMA-RO	Romania
5	Meteo Swiss	MS	Switzerland
6	Deutscher Wetterdienst	DWD	Germany
7	Swedish Meteorological and Hydrological Institute	SMHI	Sweden
8	University of East Anglia (Climatic Research Unit)	UEA	United Kingdom
9	Météo France	MF	France

# Table of contents

Abstract.....	3
1. Scientific and/or technical quality, relevance to the topics addressed by the call .....	4
1.1 Concept and objectives .....	4
1.2 Progress beyond the state-of-the-art.....	12
1.3 S/T methodology and associated work plan.....	22
2. Implementation .....	51
2.1 Management structure and procedures .....	51
2.2 Individual participants .....	53
2.3 Consortium as a whole .....	66
2.4 Resources to be committed .....	68
3. Impact.....	71
3.1 Expected impacts listed in the work programme.....	71
3.2 Dissemination and/or exploitation of project results, and management of intellectual property .....	76
4. Ethical issues.....	78
5. Consideration of gender aspects.....	78
References .....	80
Appendix A: Acronym list .....	85
Appendix B: Support letters .....	90

## Abstract

EURO<sub>4</sub>M will develop the capacity for, and deliver the best possible and most complete (gridded) climate change time series and monitoring services covering all of Europe. These will describe the evolution of the Earth system components by seamlessly combining two different but complementary approaches: regional observation datasets of Essential Climate Variables (ECVs) on the one hand and model based regional reanalysis on the other.

The project will extend, in a cost effective manner, European capacity to systematically monitor climate variability and change (including extremes) on a range of space and time scales. EURO<sub>4</sub>M will reach out with innovative and integrated data products and climate change services to policy-makers, researchers, planners and citizens at European, national and local levels. This will directly address the needs of, for instance, the European Environment Agency for their environmental assessment reports - and even provide online reporting during emerging extreme events.

As the primary source of timely, targeted and reliable information about the state of the climate in Europe, the suggested collaborative project is an important building block for GMES. The project will integrate and extend core services activities on ECVs, specifically developing the capacity required for state-of-the-art user-oriented products for monitoring of climate change. EURO<sub>4</sub>M has the potential to evolve into a future GMES service on climate change monitoring that is fully complimentary and supporting the existing operational services.

# 1. Scientific and/or technical quality, relevance to the topics addressed by the call

## 1.1 Concept and objectives

### *Concept*

As the primary source of timely, targeted and reliable information about the state of the climate in Europe, the suggested collaborative project is an important building block for the Global Monitoring for Environment and Security (GMES) initiative. No other coordinated contribution for this area exists or is currently planned. The current GMES services, which have already entered into their pre-operational phase, are not designed to provide climate change monitoring information nor reports about high impact weather and climate extremes placed in an historical context. For example, the Atmosphere service is mainly directed towards air quality and focuses on the shorter time scales. Also, the *in situ* component of GMES at present does not fully address meteorological observations. The coordination action for *in situ* data indicated in the Work Programme (p30) alone will not result in comprehensive pan-European climate datasets at a useful level of aggregation and processing.




Figure 1.1a. Observed global and European annual average temperature deviations, 1850–2007 (EEA–JRC–WHO, 2008; Figure 5.2). The figure indicates that Europe warms more rapidly than the globe (+0.74°C over the last 100 years). The annual deviations are relative to the period 1850–1899 to better monitor the EU objective not to exceed 2 °C above pre-industrial values. Over Europe average annual temperatures during the real pre-industrial period (1750–1799) were very similar to those during 1850–1899. Source: UEA/Climate Research Unit ([www.cru.uea.ac.uk/cru/data/temperature/](http://www.cru.uea.ac.uk/cru/data/temperature/)).

Climate change is the societal benefit area of the Group on Earth Observations (GEO; see Appendix A for a list of acronyms) that lacks, and urgently needs, an integrated and coordinated approach with a focus, in particular, on climate information for the multi-decadal time scales that are most relevant for adaptation. The other societal benefit areas (water, natural and human-induced disasters, environment and health, energy, ecosystem services, agriculture and desertification, biodiversity) are already reasonably well covered in the existing data information systems, such as the European Environment Information and Observation NETwork (EIONET) and the Shared Environmental Information System (SEIS). Integrated long-term and high-quality datasets of climate change information (in particular extremes) in terms of atmospheric Essential Climate Variables (ECVs) are typically missing (see [dataservice.eea.europa.eu](https://dataservice.eea.europa.eu)).

This situation is limiting the response strategies to adapt to climate variability and change at the regional, sub-regional and national scales. In particular, information on changes in weather and climate extremes is crucial, especially as the driving force for the impact work in all GMES services and GEO areas. If the relevant climate change information is not made available, then these services and areas will not be able to be successful. Due to a lack of coherent information on weather and climate extremes many GMES services and GEO areas base their work on changes in mean climate only. However, it is generally accepted that the impacts of climate change are caused primarily by changes in variability and extremes, rather than changes in the mean climate. For adaptation strategies, the longer (multi-decadal) time scales are particularly relevant, because nearly all infrastructure design relies on assessment of probabilities of extremes with return periods of  $\geq 50$  years. These assessments should take into account that the climate is non stationary because of climate change. It is the longer time scale that is needed for governments to implement their climate change action plans.

The members of this project's consortium are currently the main source of climate change time series and monitoring information for governments, policy-makers and the general public across Europe. They are frequently approached by the European Environment Agency (EEA), Joint Research Centre (JRC), World Meteorological Organization (WMO) and World Health Organization (WHO) to contribute this information to environmental assessment reports. The EURO4M participants **MO** and **UEA** collectively provide information on European temperature change over the past decades based on their global datasets (Figure 1.1a), which are also prime inputs to the reports of the Intergovernmental Panel on Climate Change (IPCC) and to the Conference of the Parties to the United Nations Framework Convention on Climate Change (UNFCCC). **KNMI** contributes up-to-date information about trends in extremes (Figure 1.1b) by linking the historical data archives of more than 40 meteorological services and universities in Europe through the European Climate Assessment & Dataset project (ECA&D; a forthcoming WMO Regional Climate Centre). **DWD** operates the Global Precipitation Climatology Centre (GPCC), which provides global analysis of precipitation on the earth's land surface based on *in situ* rain gauge data.

All these activities have an *ad hoc* character and limited spatial (horizontal and vertical) resolution for the longer time scales. Also, the potential of data assimilation and reanalysis for climate change monitoring are not yet fully exploited. **MO**, **SMHI** and **MF** are at the forefront of data assimilation developments in Europe, but no regional reanalysis for Europe is currently available that improves on the existing global reanalyses (particularly ERA-40 at the European Centre for Medium-Range Weather Forecasts ECMWF). **URV**, **NMA-RO** and **MS** have contributed significantly to improved observational datasets for sub-regions, but these activities have not yet been integrated

within the European context. DWD coordinates the Satellite Application Facility on Climate Monitoring (CM-SAF) of EUMETSAT, which aims at the provision of satellite-derived geophysical parameter datasets suitable for climate monitoring. These datasets need integration too.

**In summary, for GMES to become a success, the situation of fragmentation and scarcity of long-term climate change monitoring information in Europe needs to change. There is the vast task of integrating national observing systems, existing global and European observation datasets, satellite-derived datasets and reanalyses into GMES** (see Butler, 2007). This is needed to fill the gap for surface climatological data and information which, at present, is clearly visible in all environmental assessments. In the words of our third party participant EEA (see Section 2.3): “Everybody is expecting that weather and climate data is simply available (according to their experience having weather forecasts for every location on every day), but for historical data on climate extremes this is clearly not the case”.




Figure 1.1b. Changes in the contribution of heavy rainfall to total precipitation 1961–2006 (EEA-JRC-WHO, 2008; Map 5.9). The figure indicates that the proportion of rainfall from heavy falls increases over most areas. Source: The E-Obs dataset from the EU-FP6 project ENSEMBLES ([www.ensembles-eu.org](http://www.ensembles-eu.org)) and the data providers in the ECA&D project ([eca.knmi.nl](http://eca.knmi.nl)).

## Objectives

The overall goal of EURO<sub>4</sub>M is to develop the capacity for, and deliver the best possible and most complete (gridded) climate change time series and monitoring services covering all of Europe. These will enable adequate descriptions of the status and evolution of the Earth system components.

Specifically, the objectives of EURO<sub>4</sub>M are to:

1. **generate time series of observation datasets and reanalyses of past observational data;**
  - build on and integrate existing European *in situ* and satellite datasets, bearing in mind global connectivity, interoperability and data sharing;
  - develop the capacity for climate quality dynamic reanalysis that optimally integrates the widest possible range of *in situ* and satellite data;
  - demonstrate the capability of regional reanalysis and multi-staged downscaling with increased levels of accuracy;
  - deliver demonstration regional reanalysis for parts of the past 20 years;
  - reduce gaps and deficiencies in European monitoring capacity through a better exploitation of existing atmospheric observations and data exchange;
  - build on the strong synergy the participants have with other major climate monitoring centres worldwide, in particular with centres involved in global reanalysis.
2. **produce innovative and integrated high-quality data products for research and applications sector users;**
  - produce multi-purpose products and information to assist climate change research to incorporate the monitored ECVs;
  - provide reliable, up-to-date scientific input (especially through the IPCC) for the implementation of European and international policies and strategies on the environment and society, including the EU climate adaptation strategy;
  - provide online reporting during emerging extreme events.
3. **reach out with data products and climate change services to the user community, stakeholders, policy-makers, and general public;**
  - demonstrate the climate change services to policy-makers, researchers, planners and citizens at European, national and local levels;
  - hold frequent dialogues and interactions with a wide range of end-users to achieve a better understanding of information needs and formats;
  - make the data and information readily accessible to users with full consideration of the appropriate level of aggregation and standardization.
4. **evolve into a future GMES service on climate change monitoring that is fully complimentary and supporting the existing core services.**
  - integrate and extend core GMES services activities on ECVs, specifically developing the capacity required for user-oriented multi-purpose products for monitoring of climate change;
  - link to existing GMES services, and especially those on marine, land and atmosphere monitoring, which include – or will include in the near future – a global component by design;
  - stimulate the GMES downstream sector;
  - demonstrate and strengthen the European leadership in long-term monitoring of climate change.

These objectives will be achieved over a 4 yr period in 4 major Work Packages (WPs) detailed in Section 1.3. Together, they comprehensively address the scientific, technical and wider societal and policy objectives of the Sub-activity SPA.2009.1.1.02 “Monitoring of climate change issues (extending core service activities)”.

### **Scope**

**Variables:** EURO4M will focus on atmospheric surface climate (air temperature, sea surface temperature, precipitation, snow cover, air pressure, surface radiation budget, wind speed and direction, water vapour) and upper-air climate (earth radiation budget, upper-air temperature, wind speed and direction, water vapour, cloud properties). These are the majority of the ECVs for the atmosphere defined for the Global Climate Observing System (GCOS) and endorsed by GEO for their System of Systems GEOSS.

**Area:** EURO4M will cover Europe in its entirety. The extent of the datasets will include Region VI as defined by WMO and Europe as defined by EEA.

**Resolution:** EURO4M will provide multi-decadal gridded datasets down to 25 km horizontal resolution (obtained by direct observations) or 3 km horizontal resolution (obtained by regional reanalysis and downscaling).

**Period:** EURO4M will include the past 20-150 years. The exact period will differ between the various data sources considered (up to 150 years for *in situ* data, 30 years for satellite data, and, at this stage, parts of the last 20 years for the regional reanalysis).

In line with the GMES goals, EURO4M will not stop when the raw (observation and reanalysis) data have been made available. The project will also engage in developing the multi-purpose products and services at the appropriate level of aggregation and processing to respond to a wide range of users and downstream services.

### **Key scientific question**

The key scientific question EURO4M addresses is: “**How can we improve monitoring to help us better understand and predict climate change, extremes and weather related hazards, so that society can respond in the best possible way?**”

In order to place recent changes, fluctuations and extremes in their long-term perspective, optimal usage of all conventional (*in situ*) and satellite sources will be required. EURO4M will achieve this by seamless integration of data from the satellite era (back to the 1970s) with early instrumental observations (back to the mid-19th century). A vital approach to integrating these quantities is for EURO4M to explore the use of new regional reanalysis as well as existing global reanalyses, especially ERA-40 (Uppala *et al.*, 2005).

The focus on extremes and weather related hazards requires more detail than is strictly necessary for GCOS global climate monitoring, because as a potential future GMES service on climate change monitoring, we need to fully satisfy the information needs for sub-regional and local adaptation measures. It is at those scales where climate and weather impacts are most strongly experienced. High impact extreme events range from those of short duration (such as heavy rain and associated flooding, and windstorms) to those which extend over several days (heat waves, atmospheric pollution), several months or perhaps even years (drought). They also range from continental-to-local spatial scales.

The science included in EURO4M (in particular in WPs 1 and 2; see Section 1.3) ensures that the overall goal of developing the capacity for, and delivering the best possible and most complete (gridded) climate change time series and monitoring services



for Europe will be achieved at a high scientific level. We combine two different but complementary approaches: regional observation datasets of ECVs on the one hand and model based regional reanalysis on the other. Choosing direct observations alone would have left the potential of data assimilation unexplored; choosing reanalysis only would have run the risk of missing some important aspects in the observations and might have compromised the quality of the final monitoring products and services. We are convinced that integrated climate change monitoring products of state-of-the-art quality are needed, because most of the work in downstream GMES services and GEO areas will be based on this information.

### ***Role of reanalysis***

A reanalysis of the past atmospheric state can be obtained by combining observational datasets with a comprehensive Numerical Weather Prediction (NWP) model. Using modern data assimilation methods, a complete estimate of the atmospheric state is computed that is both dynamically consistent and optimally close to the observations. The great benefit of a reanalysis is that it provides a complete picture of the atmosphere covering the whole of the three-dimensional domain, also for the ECVs and parameters which are not routinely monitored by observations.

In a reanalysis (or rather re-assimilation) also, the consistency in time is ensured through the forecast model used in the data assimilation cycles. It carries the information from cycle to cycle with dynamical changes in order to give a best background before the observations are analysed. This 4-dimensional dynamical consistency is strong both in 3D and 4D assimilation algorithms. In 4D variational assimilation (4D-VAR) the consistency is even stronger within each time window. The dynamical model trajectory is fitted as closely as possible (as given by model and observation statistics) to all the observations within the time window of 4D-VAR (6-12 hours). It means that also atmospheric tendencies are well assimilated and the resulting analysis fields provide better estimates of moist variables like precipitation rates and cloud cover compared with what 3D methods give.

The availability of forecast model fields in the assimilation cycles provides a whole range of diagnostic quantities like radiative fluxes and accumulated precipitation where there are no observations. The accuracy of these is of course limited by the quality of the forecast model, but by basing reanalysis systems on state-of-the-art operational NWP models, we benefit from past careful validation and model development in the operational centres. Experience from global reanalyses has shown that there is a potential in the use of such model-derived quantities.

The benefit of complete multi-decadal sets of ECV products motivates EEA to investigate the use of reanalysis for future assessment studies and explore possibilities for a regional analysis in Europe (the EURRA project). However, current reanalysis products are not yet reliable enough to satisfy user needs entirely, because the basic raw observations and measurements are not consistently of climate quality. Problems arise partly because the assimilating numerical model is inevitably biased to some degree in its representation of the climate system. If observations are abundant and unbiased, they can correct the biases in background forecasts when assimilated (Simmons *et al.*, 2004). In reality, however, observational coverage varies over time, observations are themselves prone to bias, either instrumental or through not being representative of their wider surroundings, and these observational biases can change over time. This introduces trends and low-frequency variations in analyses that are mixed with the true climatic signals (see Figure 1.1c).




Figure 1.1c. Changes in summer (JJA) rainfall since 1961 estimated from different sources:  
 a) the E-Obs dataset based on station observations (see Figure 1.1b);  
 b) UKCIP08 (Jenkins *et al.*, 2007);  
 c) ERA-40 reanalysis (ECMWF)  
 The difference in trends over the UK illustrates the biases in existing global reanalysis products.

Moreover, the limited spatial resolution of current models that are affordable for global reanalysis inhibits the applicability of products for the study of local climate change and extremes.

Progress in the provision of reanalyses suitable for climate monitoring requires improvements in data assimilations systems, both in assimilating models and the analysis methods used to incorporate observational information, and improved recovery, quality control and bias correction (or homogenization) of past observations. Assessment of existing products through comparison either with high quality regional datasets or alternative global ECV products is a further important activity needed both to guide potential users as to the applicability of the products they are using and to guide where the emphasis needs to be placed for the improvement of the production systems.

**EURO4M will enhance the quality of regional reanalysis methods and deliver different levels of advanced regional reanalysis datasets for Europe.** Some of these datasets will cover most of the last 20 years, whereas other more advanced systems will be

demonstrated for a few years. There will be improvement of data assimilation systems, in resolution, and in the use of precipitation and surface data. Selected products will be evaluated in detail, comparing also with existing products from global reanalyses.

**In order to further enhance the horizontal resolution and detailed structures of the regional reanalyses, EURO4M will also perform 2-dimensional downscaling** at a considerably higher resolution than is possible with the 3D (or 4D) methods. The 2D analysis methods still combine observations with background fields from the 3D regional system, but the analysis methods are relaxed from fulfilling all the 3-dimensional balance constraints that are required for NWP. In the 2D analyses one employs a lot of regional variation given by observational statistics and by physiographic factors (land-sea and orography, mainly). In this way it is possible to analyse fine scale structures given by observations and closer to the observations than the full 3D systems can do.

We will also prepare bias-adjusted observed data inputs to climate quality reanalyses in close cooperation with ECMWF. We will contribute to the improved capability for reanalysis needed to ensure that the next generation of longer-term global and regional reanalysis is as well prepared as possible to meet the needs of climate monitoring. Our third party participant ECMWF (see Section 2.3) strongly supports this activity and will collaborate by providing the current ERA-Interim reanalysis, the raw observations and by exchanging ideas on the formats of observation files, monitoring techniques and any other possible commonalities between the regional reanalysis in EURO4M and the global reanalysis.

### ***Role of satellite data***

Satellite-derived products will supplement the reanalysis data and ground measurements. Over the oceans and sparsely populated areas satellite data are often the only data source. The quality of cloud information, the surface radiation budget, water vapour and precipitation can be significantly increased by use of satellite data within the integrated data products. The top of the atmosphere radiation budget can be only observed by satellites.

Satellite products (e.g. water vapour, cloud properties, sea ice and shortwave radiation) together with ground-based measurements will also be used to verify the reanalysis data. Biases in the radiances of satellite sensors, e.g. due to missing inter-calibration, lead to biases or breaks of homogeneity in reanalysis data as the models themselves cannot absolutely correct biases in grossly inhomogeneous input data. Satellite radiances are already assimilated into reanalysis datasets. In order to improve the homogeneity of reanalysis data, EURO4M will support satellite inter-calibration activities. The sensitivity of reanalysis ECVs for inhomogeneities and biases in satellite radiances will be evaluated.

Satellite data will be used together with reanalysis data to provide a long-term record of the surface radiation budget. Climate indices derived from satellite-based products are of value for monitoring of specific climate impacts (e.g. droughts in Wang and Qu, 2007). Additionally, the commercial solar energy market demonstrates the suitability of satellite data for climate applications. E.g. satellite-based solar irradiance data are used for the efficient planning and monitoring of solar energy systems (Hammer *et al.* 2003; Drews *et al.* 2008).

The product palette of the CM-SAF has been evaluated to be suitable for monitoring of climate change (Schultz *et al.*, 2008). However, **the overall capacity of satellite-derived data for climate monitoring has by no means been explored satisfactorily. EURO4M will**

**integrate the EUMETSAT-SAF products and methods for climate monitoring.** In this respect, reprocessing of existing satellite products in order to derive homogeneous long-term datasets is an important effort. At the same time, there is scope for merging these products with shorter-term but possibly more advanced datasets.

### ***Climate Information Bulletins and Climate Liaison Team***

Whereas the raw data from EURO4M will be distributed mainly through existing systems and the scientific results through peer-reviewed scientific and technical journals, **the EURO4M multi-purpose products will be disseminated through regularly issued Climate Information Bulletins (CIBs).** These will focus on user groups and downstream services that do not have the required expertise and knowledge to access and process the Terabytes of raw observation data or reanalysis data. CIBs will provide simple, effective and timely knowledge abstractions from EURO4M data and activities. The CIBs will be flexible and optimal products that will be responsive to current environmental and climatic events, extremes and also user needs.

**Within EURO4M a Climate Liaison Team (CLT) will actively solicit user requirements and feedback.** Through their mediation, the multi-purpose results of EURO4M will feedback directly into applications and impact assessments relevant to European societal and community needs. The CLT will also train users in the handling of EURO4M products and services through workshops, e-learning modules, podcasts and user guides covering different skill levels. In this way, EURO4M will ensure stronger links between European applications studies and the outputs from traditional and advanced climate monitoring systems.

The existing core GMES services and the downstream GMES services, which are currently planned and positioned between the multi-purpose core services and the individual user clients, will be actively involved in the CLT. This will ensure that the end-users can and will take full benefit of the wide range of products and services EURO4M will develop. The CLT will build on existing national experience and related ERA-Net activities, such as the ERA-Net: Climate Impact Research Coordination within a Larger Europe (CIRCLE; see Section 3.1 and Kabat and Vellinga, 2005). The lessons learned from the process for establishing user needs that has been used for the selection of fast track and pilot services will be considered too. The CLT enables that, already in an early stage, users are involved in the decision-making process in a structured way. This is in line with the user-driven objective of GMES (EC, 2008).

## **1.2 Progress beyond the state-of-the-art**

### ***Innovation-related activities***

Never before has the pan-European integration of atmospheric observations from ground-based sources, satellite sources and reanalysis been optimized in such a comprehensive way for long-term climate monitoring and adaptation policy support. Remotely sensed data from satellites deliver near-real time estimates of key parameters worldwide. However, they must be complemented by *in situ* data, even though these are often much more sparsely and irregularly distributed in time and space. The *in situ* data are indispensable for calibration and validation of satellite data. Also they give information representative of a finer spatial scale, often with higher precision, which complements satellite information which is coarser, but has wider coverage.

The problems with estimates of lower tropospheric temperatures from satellite-based Microwave Sounding Unit (MSU) radiances since 1978, and their subsequent comparisons with sonde-based and surface temperature data, indicate that datasets need to be considered together rather than in isolation. In contrast to satellite data, *in situ* measurements have often been acquired and handled by a wide variety of laboratories and institutions for the purposes of specific research programmes and are not managed in a unified way.

### ***Essential Climate Variables (ECVs) state-of-the-art***

The Second Adequacy Report of GCOS (GCOS, 2003) developed the concept of Essential Climate Variables (ECVs) encompassing the atmospheric, oceanic and terrestrial domains. The concept forms part of the GCOS Implementation Plan (GCOS, 2004), which for these three domains has been endorsed by GEO in its GEOSS work plan (GEO, 2007). Note that the GCOS Implementation Plan will be updated, starting from a meeting on 2-5 February 2009, in which EURO4M partners will be involved.

GCOS named the ECVs in a somewhat *ad hoc* way. Some are very specific physical quantities, e.g. surface temperature and precipitation, while others are more vague and generic, e.g. cloud properties and minor greenhouse gases, encompassing a number of measurements. Some can be considered at point locations, while others only really have scientific relevance as global fields.

**EURO4M will consider a selection of ECVs** from the atmospheric domain, which comprises 6 surface level, 5 upper atmosphere ECVs, and only one ECV from each of the oceanic and terrestrial domains (see Table 1.2a).

Domain	Essential Climate Variables (ECVs)
Atmospheric (over land, sea and ice)	Surface: <b>Air temperature, Precipitation, Air pressure, Surface radiation budget, Wind speed and direction, Water vapour</b> Upper-air: <b>Earth radiation budget, Upper-air temperature, Wind speed and direction, Water vapour, Cloud properties</b> Composition: Carbon dioxide, Methane, Ozone, Other long-lived greenhouse gases, Aerosol properties
Oceanic	Surface: <b>Sea-surface temperature</b> , Sea-surface salinity, Sea level, Sea state, Sea ice, Currents, Ocean colour, CO <sub>2</sub> partial pressure Sub-surface: Temperature, Salinity, Currents, Nutrients, Carbon, Ocean tracers, Phytoplankton
Terrestrial	River discharge, Water use, Ground water, Lake levels, <b>Snow cover</b> , Glaciers and ice caps, Permafrost and seasonally-frozen ground, Albedo, Land cover, Fraction of absorbed photosynthetically active radiation, Leaf area index, Biomass, Fire disturbance.

Table 1.2a. GCOS ECVs: Those that will be part of EURO4M are shown in bold.

Specific reasons for this selection of ECVs are justified in the assessment below. The general motivation is that **these ECVs form primary input to all GMES services and GEO societal benefit areas** and that most services and areas currently make suboptimal use of existing information. Often the changes in extremes that accompany the average warming are not taken into account. Also, most of the other ocean and terrestrial ECVs are part of other core GMES services. With several other projects underway that are fully dedicated to monitoring the atmospheric composition (such as CarboEurope and NitroEurope), these ECVs are not considered within EURO4M.

### ***ECV: Surface Temperature***

The best-known product for monitoring surface temperature, using a combination of land station data and *in situ* sea surface temperature observations, has been developed by the Climatic Research Unit of UEA and the MO. The current version of the monthly dataset, which extends back to 1850, is HadCRUT3 (Brohan *et al.*, 2006) but its spatial and temporal resolutions of 5° by 5° latitude/longitude and monthly timescale are rather coarse. A higher resolution version, back to 1901 at 0.5° by 0.5° spatial resolution, is available for land areas only (Mitchell and Jones, 2005) but the temporal resolution is still monthly. Sea surface temperature data is well studied (as it is a vital input to Reanalyses) and datasets such as HadISST2 and HadSST2 (Rayner *et al.*, 2006) are readily available. National Meteorological and Hydrological Services (NMHSs) have collected most of the historical station temperature data, but only a few have digitized their entire daily to sub-daily holdings. Fewer still have homogenized their long series. Some additionally charge for accessing this type of data.

Recent EU projects EMULATE (European and North Atlantic daily to MULTidecadal climate variability), ENSEMBLES (ENSEMBLE-based Predictions of Climate Changes and their Impacts), STARDEX (Statistical and Regional dynamical Downscaling of Extremes for European regions), IMPROVE (Improved Understanding of past climatic variability from early daily European instrumental sources), CIRCE (Climate change and impact research: the Mediterranean environment) and ECA&D have assembled daily series of maximum and minimum temperature, all of which will be available to EURO4M. ENSEMBLES and STARDEX have emphasised high spatial density observations, but over relatively short periods from the 1950s. EMULATE, CIRCE and ECA&D concentrated on records that extended back to the 19th century, but their databases were limited to about 200 records, and spatial gaps continue to affect Europe. ENSEMBLES has produced a daily gridded dataset at a resolution of 25 by 25 km for statistical comparisons with Regional Climate Model output (Haylock *et al.*, 2008). Higher-resolution (3 hourly) data are also digitally available from the Integrated Surface Hourly (ISH) dataset. These are discussed under the surface humidity ECV, but Integrated Surface Hourly (ISH) temperatures will also be investigated in EURO4M.

In summary, temperature is the ECV for which most digitized data are readily available, and it is also the most widely analyzed variable. Some spatial limitations affect Europe in the daily series, and there are concerns about the long-term homogeneity of the records (Wijngaard *et al.*, 2003).

### ***ECV: Precipitation***

Gridded precipitation datasets are available at 0.5°, 1° and 2.5° resolution from the WMO Global Precipitation Climatology Centre (GPCC) operated by DWD. These are at the monthly timescale, extend back to 1951, have global coverage (land-surface), and use

between 10,000 and 40,000 individual station series (Rudolf and Schneider, 2005, Beck *et al.*, 2005). Since the late 1970s, it has been possible to incorporate satellite precipitation estimates, although these raise a myriad of issues regarding the types of satellite sensors used and the algorithms which convert sensor measurements into surface precipitation rates. Accordingly, global datasets of precipitation have been constructed by the Global Precipitation Climatology Project (GPCP) and NOAA's Climate Prediction Center Merged Analysis of Precipitation, both being available at 2.5° resolution. Blending of the ground-based GPCP precipitation dataset with the satellite-based HOAPS dataset (Hamburgs Ocean Atmosphere Parameters from Satellite; Grassl *et al.*, 2000; Bauer and Schuessel, 1993) lead to a global precipitation dataset covering both land surface and ocean (Figure 1.2a).




Figure 1.2a: Annual average precipitation amount derived by blending GPCC with HOAPS-3. DWD will generate a blend of HOAPS-4 and GPCC, which will comprise 20y of global precipitation over land (GPCC) and ocean (HOAPS).

Slightly more digitized daily records are available for precipitation than temperature across Europe, but given the much greater spatial variability of precipitation, a much greater density is essential. This is particularly the case for mountain regions (Frei and Schär 1998). ENSEMBLES has produced a daily gridded dataset at a resolution of 25 by 25 km for statistical comparisons with Regional Climate Model output (Haylock *et al.*, 2008). In specific regions and projects (e.g. the Alps, the Baltic Sea Experiment (BALTEX), the European Land Data Assimilation System to predict Floods and Droughts (ELDAS) and for many individual countries), much denser networks have been used in regional and national studies. Gridded versions of these data will be used to assess the accuracy of interpolation in ENSEMBLES (extending the initial work reported by Hofstra

*et al.*, 2008). As for temperature, CIRCE aims at collecting, updating and homogenising daily precipitation station series from the larger Mediterranean area covering the last decades to 100 years. The WMO MEDARE initiative aims at pushing this limit even further back in time and recover additional series from this region.

In summary, precipitation data need to be more spatially extensive than temperature to achieve a given level of accuracy in regional-scale estimations. There are again limitations with the availability of digitized data across Europe, and although there are fewer concerns about homogeneity, this may be because the density of networks is inadequate to address this issue (Auer *et al.*, 2005). EURO4M will have access to daily station series as well as gridded products from GPCP, EMULATE, ENSEMBLES, CIRCE and MEDARE (see Section 1.3).

### ***ECV: Atmospheric Air Pressure***

Most analyses of atmospheric circulation in the scientific literature during the last few years make use of reanalyses (ERA-40 or NCEP/NCAR). However, efforts to examine circulation patterns over longer time scales have focused on observational datasets of atmospheric pressure. Foremost amongst these, using both global terrestrial ISPD and marine ICOADS data at monthly timescales back to 1850 (and at 5° by 5° spatial resolution), is the Hadley Centre mean Sea Level Pressure dataset, Version 2 (HadSLP2) compiled by MO (Allan and Ansell, 2006). The same institution led the EMULATE project in its development of a daily (24 hour mean) mean sea level pressure (MSLP) product from 1850, also including both terrestrial and marine ICOADS data at the same resolution (EMSLP), but only for the North Atlantic-European region 25-70°N by 70°W-50°E (Ansell *et al.* 2006). EMULATE made extensive use of earlier EU-funded projects that digitized long daily pressure series, such as WASA (Waves and Storms in the North Atlantic) and IMPROVE. The basic terrestrial and island station pressure data in the ISPD and the marine measurements from ICOADS, that are the basis of the above gridded datasets, are being expanded and improved through the Atmospheric Circulation Reconstructions over the Earth (ACRE) initiative (led by MO) and will be available for EURO4M. The importance of atmospheric air pressure for climate change monitoring is illustrated by Figure 1.2b from IPCC-WGI-AR4. Here, SMHI have used station data to indicate evidence for changes in extratropical cyclone activity (storminess).

Efforts are underway at MO via the ACRE initiative and its links to the 20th Century Reanalysis Project of its US collaborators, to develop the combined ISPD and ICOADS global sub-daily MSLP databases for climate quality surface-observations-only reanalyses extending back to 1891. Additional ACRE-facilitated historical reanalyses using only surface observations are planned to extend these reconstructions back over the last 150 years globally and for the North Atlantic-European region to the mid-18<sup>th</sup> century. The development of the combined ISPD and ICOADS sub-daily MSLP dataset needed for such activities will be closely linked to EURO4M activities in WP1. All of these new MSLP data activities are structurally, but not financially, supported by GCOS through the AOPC/OOPC Surface Pressure Working Group.

Station pressure data are rarely analyzed in isolation (except for the WASA work with pressure triangles) and most studies derive MSLP fields. For Europe, these fields are generally adequate, but will be improved and extended back in time over both the land and oceans by the ACRE project. Improvements are required for sub-daily fields and in marine data in some parts of the world, and new surface-observations-only reanalyses should begin to address these deficiencies.






Figure 1.2b: Storm index for the British Isles, North Sea and Norwegian Sea, 1881 to 2004. Blue circles are 95th percentiles and red crosses 99th percentiles of standardised geostrophic winds averaged over 10 sets of triangles of stations. The smoothed curves are a decadal filter (updated by SMHI from Alexandersson *et al.*, 2000; see Figure 3.41 in IPCC-WGI-AR4).

### ***ECV: Surface and Earth Radiation***

The surface radiation budget is a fundamental component of the surface energy budget that is crucial to nearly all aspects of climate. The Baseline Surface Radiation Network (BSRN) of the World Climate Research Programme (WCRP) has established the relevant measurement techniques and is now recognized as the GCOS baseline network for surface radiation. The BSRN provides high-quality but spatially limited measurements of radiation at the surface. The GCOS action to expand the BSRN network, increasing the number of land stations and using research ships and buoys over the ocean is an important step. However, additional data are needed to obtain a sufficient spatial coverage for climate monitoring. As with most GCOS actions and plans, the initiatives and financial resources must come from the scientific community, principally through the member states and their NMHSs.

Satellite data can be used to monitor clouds and surface solar radiation quite well. Hence BSRN stations in combination with satellites are a powerful option for an appropriate monitoring of the surface radiation with high accuracy and spatial coverage. The International Satellite Cloud Climatology Project (ISCCP), the Earth Radiation Budget Experiment (ERBE), as well as the CM-SAF Climate Data Records will be used within EURO4M. The potential of merging *in situ* data with satellite-derived data will be investigated in WP1.

In addition, reanalysis data has the potential to contribute to the appropriate monitoring of radiation ECVs. Trentmann *et al.* (2008) has recently evaluated the thermal flux densities at the surface of the ERA-Interim dataset and concludes that this has a high

accuracy and can therefore be used in conjunction with satellite-based shortwave surface radiation fluxes in order to provide the surface radiation budget and its components in climate accuracy. Of course, monitoring the radiation budget at the top of the atmosphere cannot be based on surface data, but needs satellite data. WP2.4 will address the accuracy of reanalysis data by comparison with the radiation budget derived with GERB/CERES (Geostationary Earth Radiation Budget/Clouds and the Earth’s Radiant Energy System) and ERBE data.

One of the radiation budget components, the surface solar radiation, is necessary for an efficient planning and monitoring of solar energy systems (Hammer *et al.*, 2003; Drews *et al.*, 2008). Surface solar radiation is also important for the satellite-based estimation of drought and evaporation (Wang and Liang, 2008; Wang and Qu, 2007).

**ECV: Wind Speed and Direction**

This ECV is measured at many stations across Europe, but because of local effects, studies of longer-term changes in high wind speeds during storms generally make use of daily and sub-daily MSLP fields (from reanalyses or from the EMULATE and WASA projects) as these are generally more consistent through time.

Figure 1.2c illustrates the use of reanalysis data for the reconstruction of past extreme wind storm events. The violent storms that swept Western and central Europe from 25 to 28 December 1999, left more than 60 dead.




Figure 1.2c: Reconstruction of the 1999 “Christmas” storms using ERA-40 reanalysis data. This example shows the wind field of 27 December 1999 at 00 UTC. Colours indicate wind speed in Beaufort scale F (source: ECMWF).

### ***ECV: Surface Humidity***

HadCRUH is a new dataset of monthly mean surface humidity anomalies at  $5^{\circ} \times 5^{\circ}$  resolution, covering the period 1973–2003, with respect to the 1974–2003 climatology (available at [www.hadobs.org](http://www.hadobs.org); see Willett *et al.*, 2007; 2008). It is nearly global in coverage, combining land data from observing stations and marine data from ships, buoys and observing platforms. The land data are sourced from version 2 of the integrated surface hourly (ISH) dataset, supplied by the National Climatic Data Center. The marine data are sourced from release 2.1 of ICOADS (International Comprehensive Ocean-Atmosphere Dataset) for 1973–1997 and from NCEP GTS data (Global Telecommunications System data made available through NOAA’s National Centers for Environmental Prediction) for 1998–2003. HadCRUH is available in specific humidity and relative humidity forms. This dataset uses all available humidity observations (vapour pressure, relative humidity and dew point temperature). EURO4M will analyse the dataset for changes over the last 30 years. The GTS dataset will also be investigated for some of the other surface ECVs (e.g. temperature and pressure), as it will likely provide sub-daily resolution for the European domain.

### ***ECVs: Upper Air Temperature/Upper Air Humidity/Wind Speed and Direction***

Upper-air temperatures, humidities and wind speeds/directions from radiosondes are a cornerstone of the data input to most reanalyses. However, improvements in the response times and radiative shielding of instruments on the sondes mean that all upper-air temperatures and humidities are subject to major heterogeneities throughout the 50+ yr record. A significant change to ascent times took place in 1957, partly explaining why ERA-40 begins in 1958. Changes to the times of the sonde ascents can affect the long-term homogeneity of the derived series through changes in the effect of radiation on the sensors as well as through the true, though small, diurnal cycle of atmospheric temperature. Some important advances in radiosonde homogeneity have been made in recent years, e.g. Thorne *et al.* (2005a, b) and the MO gridded radiosonde temperature dataset (HadAT). At the National Climatic Data Center (NCDC, US), many of the original sonde ascents were collected or digitized in the Integrated Global Radiosonde Archive (IGRA; Durre *et al.*, 2006).

WP2 will benefit significantly from previous work using data from both HadAT (homogeneity adjustments, see Titchner *et al.*, 2008) and IGRA (additional pre-1979 ascents). Data homogenization by near neighbour-comparisons, and by reference to reanalysis feedback files (e.g. by Haimberger *et al.*, 2008), will be incorporated into improved analyses for the European-North Atlantic region.

### ***ECV: Cloud Properties***

Cloud feedback is considered to be the single most uncertain aspect of future climate projections and is responsible for much of the wide range of estimates of climate sensitivity in climate models. Therefore, it is vital to maintain long-term records of cloud properties. Satellite data, for example constructed by ISCCP, provide an appropriate basis for monitoring of cloud parameters. EURO4M will deliver a number of new cloud property datasets produced within the framework of the CM-SAF (including a 20+ yr cloud albedo dataset). Monitoring of developments within ISCCP and a liaison with AOPC will ensure that the most up-to-date information is used. A metric for evaluation of reanalysis cloud data will be developed using ISCCP, CM-SAF and cloud products derived in WP2.1.

### ***ECV: Snow cover***

This ECV is classed by GCOS as being in the terrestrial domain, but it has traditionally been measured by NMHSs. Thus across northern and high-elevation areas of Europe, there are long daily records of station-observed snow cover, many of which are digitally available. In addition to the *in situ* observations, satellites have measured snow cover since about 1966. Fields of snow cover have been quite deficient in previous reanalyses, and experiments with improved fields will be investigated within WP2. Data for this ECV will also be available through interactions between EURO4M and the EuroCryoClim project for high latitude Europe and the Arctic.

### ***New research initiatives***

**High quality, high-resolution data products from EURO4M will provide the information basis for new generations of research initiatives** (e.g. initialization and evaluation of Earth System Models, climate change detection and attribution studies) **and wider societal, industrial and environmental user applications** (e.g. climate risk, impact assessment, study of socio-economic effects of extreme climate events, reducing loss of life and property from disasters, managing the consequences of climate change, development of integrated management strategies).

EURO4M will provide a vital stimulus for the next generation of reanalyses, including those extending further back in time. Analyses of observations used in existing reanalysis products, and those that could have been used, will highlight what can be achieved if all available observations are improved and used where possible.

These new research initiatives fully comply with the recommendations for future EU integrated climate research resulting from the IPCC-AR4, which have been discussed at the international symposium “Future climate, impacts and responses” organised by the European Commission, Research Directorate-General in Brussels, 19-20 November 2007.

### ***Climate quality reanalysis***

**The data assimilation and downscaling activities in EURO4M will progress the regional reanalysis science beyond the present state-of-the-art.** The data assimilation systems are making rapid progress as they benefit from research in NWP, which has led to much improved weather-forecast skill. Also, progress is made in the correction of biases and discontinuities of the observing system that obscure the detection of climate trends. Computing resources are growing with time, which will enable future reanalyses to be calculated at finer horizontal and vertical resolutions, with immediate effects on product quality and on the relevance of products for new applications.

An example of what can be achieved is provided by the National Centers for Environmental Prediction (NCEP) North American Regional Reanalysis (NARR). This long-term, consistent, high-resolution climate dataset for the North American domain is a major improvement upon the earlier global reanalysis datasets in both resolution and accuracy (Mesinger *et al.*, 2006). At 32 km horizontal resolution, NARR covers the 25-yr period 1979–2003 and is being continued in near-real time. In particular, NARR has successfully assimilated high-quality and detailed precipitation observations into the atmospheric analysis. Consequently, the forcing to the land-surface model component of the system is more accurate than in previous reanalyses, so that NARR provides a much-improved analysis of land hydrology and land–atmosphere interaction. The overall

atmospheric circulation throughout the troposphere has been substantially improved as well.

Operational agencies responsible for reanalysis need to be kept abreast of advances in data recovery and rehabilitation, which will lead to improvements in datasets for assimilation in reanalysis. **In the long run, new “climate quality” dynamic reanalyses will then be able to provide the necessary ECVs.** This requires data from a network of radiosonde, surface-based, and satellite-based observations that are specifically pre-validated with respect to systematic biases. The models themselves cannot absolutely correct biases in grossly inhomogeneous input data. Incorrect or incomplete data with spatio-temporal inhomogeneities can be misleading in estimating climate change, in particular changes in extremes. Recovery of synoptic surface meteorological data and radiosonde upper-air data is needed to fill gaps in the observational records held by reanalysis centres. Major efforts are needed to bring together more of the available data into a coherent form suitable for climate quality reanalyses. For instance, existing reanalysis products did not use all corrected surface-based and radiosonde observations. For the early years, much better exploitation of existing and digitized high-resolution observational series can be made. EURO4M will therefore extend these data back well before 1958 (the start date of ERA-40) to serve future reanalysis prior to this time, e.g. the proposed next generation reanalysis (ERA-75) that ECMWF plans to carry out, with “proof of concept” being undertaken by the ERA-CLIM proposal for FP7 should it be successful (see Section 2.3). The project will also link with the ACRE-facilitated reanalyses to aid the development and testing of a 100+ year reanalysis using only surface observations (the 20th Century Reanalysis Project), and thus be positioned to work similarly with even longer historical climate quality reanalyses.

## 1.3 S/T methodology and associated work plan

### *Strategy of the work plan*

The scientific and technical approach of EURO4M has been designed to achieve the project goal and objectives (as stated in Section 1.1) in a realistic, measurable and specific manner within **4 years**. EURO4M will combine an integrated European climate monitoring capacity (supply) with user needs (demand). Improved data coordination and analyses will help fill gaps in the atmospheric observation system, and contribute to advanced data assimilation and reanalyses. High quality integrated ECV datasets and aggregated climate change products and services (CIBs) form the major outputs of the project.

### *Data exchange*

Many NMHSs have been made to, or found it necessary to, charge for the acquisition of such data, especially daily to sub-daily weather records. This has put some NMHSs data outside of the budgetary realms of research institutes, organisations and agencies. It may also be one of the reasons that integrated datasets of climate change information (in particular extremes) are typically missing in the existing data information systems, such as EIONET and SEIS (see [dataservice.eea.europa.eu](http://dataservice.eea.europa.eu)). No uniform data policy for climatological data exists in Europe. The Network of European Meteorological Services (EUMETNET) has struggled with this topic for many years now without reaching a

solution. Large differences continue to exist between individual countries and there is no single European counter (or portal) for climatological data.

**In EURO4M, we will respect the data policies of NMHSs and other data providing institutions. At the same time, we will also respect and try to work in the spirit of GMES and WMO:**

- GMES services, according to the Commission (EC, 2008), should be fully and openly accessible, as long as EU and Member States security interests do not suggest otherwise. This will help to promote the widest possible use and sharing of Earth observation data and information in line with the proposed SEIS and in accordance with existing legislation such as the Infrastructure for SPatial InfoRmation in Europe (INSPIRE) Directive taking into account the Global Earth Observation System of Systems (GEOSS) principles.
- WMO Resolution 40 on free exchange of data (Res40Cg-XII WMO policy and practice for the exchange of meteorological and related data and products including guidelines on relationships in commercial meteorological activities and its annex 4: definitions of terms in the practice and guidelines) states that: “As a fundamental principle of the World Meteorological Organization (WMO), and in consonance with the expanding requirements for its scientific and technical expertise, WMO commits itself to broadening and enhancing the free and unrestricted exchange of meteorological and related data and products”.

Good examples of how this will work in practice are available from the ongoing daily station data collation for the ECA&D and the ENSEMBLES project, as well as the reanalysis data from ECMWF and the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR).

### *Timing of WPs*

**The EURO4M work plan is organised as 4 interacting WPs (see Table 1.3a), each of which are broken down into sub-WPs (= Tasks). The rationale for this structure is described in the section “graphical presentation of the components”, which immediately follows the descriptions of the individual WPs. The project structure reflects that we will combine two different but complementary approaches: regional observation datasets of ECVs on the one hand (WP1) and model based regional reanalysis on the other (WP2). In this way, we retain the best of both worlds delivering the best possible and most complete climate change monitoring services for Europe (WP3).**

Gantt charts for the WPs and deliverables/milestones are shown in Figure 1.3a,b. Note that the names for the (sub) WPs and deliverables/milestones in the charts are only brief summaries of the full descriptions provided in the Tables 1.3a-d. These charts provide a general picture of the timing of the project. More detailed interdependencies between the various tasks will be described in the section following the WP descriptions of work. Important to note here is that an advanced system for multi-decadal regional reanalysis, which will use future ERA75 boundaries, will be developed in parallel to the production of a multi-decadal regional reanalysis dataset. This will guarantee that adequate regional reanalysis data can already be used within the project to demonstrate the integrated products and services.

WP gantt chart																																																			
Month:	Year 1:												Year 2:												Year 3:												Year 4:														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48			
<b>WP1 Regional observation datasets</b>																																																			
WP1.1 Gridded datasets – stations																																																			
WP1.2 Gridded datasets – remote sensing																																																			
WP1.3 Data coordination																																																			
<b>WP2 Regional reanalysis</b>																																																			
WP2.1 Advanced regional data assimilation																																																			
WP2.2 Dynamical downscaling of ERA																																																			
WP2.3 2D-mesoscale downscaling																																																			
WP2.4 Evaluation																																																			
WP2.5 Improved input data for reanalysis																																																			
<b>WP3 User oriented information/products</b>																																																			
WP3.1 Climate Indicator Bulletins (CIBs)																																																			
WP3.2 Climate Liaison Team (CLT)																																																			
<b>WP4 Project management</b>																																																			

Figure 1.3a. GANTT diagram for WPs




Figure 1.3b. GANTT diagram for Deliverables and Milestones



*Work description broken down into WPs and sub-WPs*

Table 1.3a Work Package list

Work Package No	Work Package title	Type of activity	Lead participant No	Lead participant short name	Person-months	Start month	End month
<b>WP1</b>	<b>Regional observation datasets</b>	RTD	8	UEA	235	1	48
WP1.1	<i>Gridded datasets – stations</i>	RTD	5	MS		1	48
WP1.2	<i>Gridded datasets – remote sensing</i>	RTD	6	DWD		1	48
WP1.3	<i>Data coordination</i>	RTD	3	URV		1	36
<b>WP2</b>	<b>Regional reanalysis</b>	RTD	2	MO	281	1	48
WP2.1	<i>Advanced regional data assimilation</i>	RTD	2	MO		1	48
WP2.2	<i>Dynamical downscaling of ERA</i>	RTD	7	SMHI		1	36
WP2.3	<i>2D-mesoscale downscaling</i>	RTD	9	MF		13	48
WP2.4	<i>Evaluation</i>	RTD	5	MS		25	48
WP2.5	<i>Improved input data for reanalysis</i>	RTD	8	UEA		13	42
<b>WP3</b>	<b>User-oriented information / products</b>	RTD	4	NMA-RO	107	1	48
WP3.1	<i>Climate Indicator Bulletins (CIBs)</i>	RTD	1	KNMI		19	48
WP3.2	<i>Climate Liaison Team (CLT)</i>	RTD	6	DWD		1	48
<b>WP4</b>	<b>Project management / coordination</b>	RTD	1	KNMI	22	1	48
				TOTAL	645		

**Table 1.3b Deliverables list**

Del. No.	Deliverable name	WP No.	Nature	Dissemination level	Delivery date
DI.1	Gridded high-resolution daily precipitation dataset for the Alpine region	1	D	PU	Month 48
DI.2	Analysis of daily to decadal variations of precipitation in the Alps	1	R	PU	Month 48
DI.3	European window of the GPCP dataset available for the EURO4M	1	D	PU	Month 6
DI.4	Extended and updated ENSEMBLES gridded daily dataset for Europe (E-Obs, incl. snow cover)	1	D	PU	Month 6, 12, 18, etc.
DI.5	Paper on extended and updated E-Obs dataset	1	R	PU	Month 42
DI.6	New UEA (CRU) data products for Potential Evapotranspiration (PET) and PDSI	1	D	PU	Month 48
DI.7	Gridded 20 yr dataset of surface solar radiation, cloud albedo, cloud fraction and surface albedo derived from MVIRI (SEVIRI) using the Heliosat method (5x5km resolution)	1	D	PU	Month 42
DI.8	Integrated gridded dataset of HOAPS and GPCP precipitation covering 20 years (1986-2006) at 0.5° x 0.5°	1	D	PU	Month 42
DI.9	Gridded dataset of water vapour based on ATOVS instrument onboard of NOAA satellites	1	D	PU	Month 42
DI.10	High-resolution datasets of precipitation, SSI and cloud properties for the MSG period (2005-present)	1	D	PU	Month 24
DI.11	Max. and min. temperature and sunshine datasets based on Land-SAF	1	D	PU	Month 36
DI.12	Update and gap-filling of pressure, temperature and precipitation data for the Mediterranean	1	D	PU	Month 30

D1.13	Merged climate dataset for the Mediterranean	1	D	PU	Month 36
D1.14	Proposal for additional data rescue activities required for data that have not yet been digitised	1	R	PU	Month 12
D2.1	New state-of-the-art NAE-based regional atmospheric data assimilation reanalysis system	2	O	PU	Month 36
D2.2	Report describing the new system in D2.1	2	R	PU	Month 48
D2.3	HIRLAM-based reanalysis dataset at 25 km for most of the past 20 yr period	2	D	PU	Month 36
D2.4	Improved MESAN downscaled dataset at 3-12 km resolution for most of the past 20 yr period	2	D	PU	Month 36
D2.5	New MESAN-SAFRAN downscaling system	2	O	PU	Month 36
D2.6	Report describing the new system in D2.5	2	O	PU	Month 48
D2.7	Evaluation of Alpine precipitation in the regional reanalysis	2	R	PU	Month 48
D2.8	Comparison of the different reanalysis downscaling products for precipitation	2	R	PU	Month 48
D2.9	Comparison of the regional reanalyses products with newly developed and existing state-of-the-art systems	2	R	PU	Month 48
D2.10	Comparison of existing ERAMESAN with SAFRAN downscaling	2	R	PU	Month 12
D2.11	Evaluation of the newly developed MESAN-SAFRAN system, in particular for snow	2	R	PU	Month 36
D2.12	Evaluation of reanalysis surface solar radiation, cloud properties, top of atmosphere albedo and water vapour using CM-SAF and WP1.2 products	2	R	PU	Month 42
D2.13	Synthesis report for all evaluation activities in EURO4M	2	R	PU	Month 48

D2.14	Analysis of the improvement of capability for reanalyses through better basic input data	2	R	PU	Month 42
D2.15	Access to <i>in situ</i> datasets not used in existing reanalysis	2	R	PU	Month 48
D3.1	Climate Indicator Bulletins (CIBs) as sound user-oriented multi-purpose products for future GMES services	3	O	PU	Month 24,36,48
D3.2	CIB for the first emerging extreme event within the course of the project	3	O	PU	Unknown
D3.3	Climate Liaison Team (CLT)	3	O	PU	Month 12
D3.4	Meta user-requirement document for the intersection of user needs across the range of user communities	3	R	PU	Month 18
D3.5	Data storage and dissemination document describing the INSPIRE-compliant infrastructure	3	R	PU	Month 48
D3.6	Two video podcasts, E-learning and other training material demonstrating EURO4M products and services	3	O	PU	Month 42
D3.7	Two workshops to support user feedback and dissemination of EURO4M products and services	3	O	PU	Month 18,48
D3.8	User guide describing the EURO4M products and services	3	R	PU	Month 48
D4.1	Kick-off meeting and first meeting of the MB	4	O	PU	Month 3
D4.2	Follow-up meetings of the MB	4	O	PU	Month 6,12,...
D4.3	Annual General Assemblies	4	O	PU	Month 12,24,...
D4.4	EURO4M website	4	O	PU	Month 1
D4.5	EURO4M brochure and logo	4	O	PU	Month 1
D4.6	Activity reports and management reports for the Commission	4	R	PU	Month 12,24,...

D4.7	Detailed implementation plan for the next 18 months for the Commission	4	R	PU	Month 1,2,24,...
D4.8	Plan for sustaining and maintaining the EURO4M CIBs as GMES climate change monitoring service	4	R	PU	Month 48
D4.9	Major final EURO4M conference showing the major project results and the way ahead	4	O	PU	Month 48

**Table 1.3c Milestones list**

<b>Milestone number</b>	<b>Milestone name</b>	<b>Work package(s) involved</b>	<b>Expected date</b>	<b>Means of verification</b>
M1.1	Gridded Alpine dataset	1	Month 48	Dataset available
M1.2	E-Obs dataset extended and updated (incl. snow cover)	1	Month 18	First version dataset available
M1.3	18 yr satellite-based dataset of radiation and clouds	1	Month 42	First version available
M1.4	Merged climate dataset for the Mediterranean	1	Month 36	Dataset available
M2.1	New state-of-the-art NAE-based regional atmospheric data assimilation system	2	Month 36	System available
M2.2	HIRLAM-based reanalysis dataset at 25 km	2	Month 36	Dataset available
M2.3	New MESAN-SAFRAN downscaling system	2	Month 36	System available
M2.4	Evaluation synthesis report	2	Month 48	Report published
M3.1	First Climate Indicator Bulletin (CIB)	3	Month 24	CIB published
M3.2	Climate Liaison Team	3	Month 12	Team established
M4.1	First General Assembly	4	Month 12	Meeting report published

Table 1.3d Work package description – part 1

Work Package number	WP 1		Start date or starting event:				Month 1	
Work Package title	Regional observation datasets							
Activity Type								
Participant number	1	2	3	4	5	6	7	
Participant short name	KNMI	MO	URV	NMA-RO	MS	DWD	SMHI	
Person-months per participant	32	24	42	36	36	16		
Participant number	8	9						
Participant short name	UEA	MF						
Person-months per participant	31	18						
<p><b>Objectives</b></p> <ul style="list-style-type: none"> <li>• further develop station-based gridded datasets for Europe in its entirety and selected sub-regions</li> <li>• assess the capacity of additional satellite-derived data for climate monitoring (linked to the EUMETSAT-SAFs), examining the parameters to be monitored, drawbacks and strengths of the methods and approaches used, the capacity for monitoring patterns of anomalies in time and across space, the capacity for trend analysis, and the capacity for the appropriate merging of satellite data with ground-truth (i.e. <i>in situ</i> based observations);</li> <li>• provide gridded satellite-based datasets for ECVs, in particular those part of the energy-water cycle, such as water vapour, components of the radiation budget at the surface and cloud albedo;</li> <li>• work with existing data recovery, rescue, imaging and digitisation activities to improve and temporally extend the ECV databases for the European region (linking with WG 1 of the international ACRE initiative plus EuroCryoClim in the Arctic, EUMETNET-ECSN for Europe, and MEDARE for the Mediterranean), in order to coordinate and make accessible currently available historical climate data;</li> <li>• conduct a comprehensive European climate data survey to identify the specific strengths and deficiencies in both the current atmospheric observation system and historical data availability (linking with WG 1 of the international ACRE initiative), focusing on the GCOS ECVs;</li> <li>• foster coordinated approaches to standards-based data management across climate-related data centres, including developing best-practice approaches for metadata, data, and services.</li> </ul> <p><b>Sub-WPs</b> (with lead participant in brackets)</p> <p>WP1.1: <i>Developing gridded datasets based on long-term station series (MS)</i></p> <p>WP1.2: <i>Developing gridded datasets based on long-term remote sensing data (DWD)</i></p> <p>WP1.3: <i>Data coordination and access to national archives (URV)</i></p>								

## Description of work

### WP1.1 Developing gridded datasets based on long-term station series (MS)

Although the best data products make best use of *in situ* and remotely-sensed information, it is important to develop some products from one or the other type of information. Provided trends and spatial patterns of change agree, the scientific certainty gained from totally independent measurements is immense. This WP will further develop station-based gridded datasets for Europe in its entirety and selected sub-regions. These will be used in WP2 for downscaling and WP3 for integrated products and services.

**MS** will develop a gridded dataset of daily precipitation for the entire Alpine region, for the period 1971 to 2007 (or more recent). Use will be made of all accessible rain-gauge observations, including the high-resolution climatological and hydrological networks of Alpine countries. Network density is a critical factor for accurate quantification of precipitation in complex terrain. This work-task will be based on the extension of an existing pan-Alpine precipitation dataset (Frei and Schär, 1998), which was collated in previous activities. The combination of high-density observations with state-of-the-art spatial analysis and modern data quality procedures aims at an accurate representation of mesoscale patterns. Particular consideration will be given to avoid spurious long-term trends due to variable station density. This knowledge is also important for the other gridded datasets in this WP. Within EURO4M, the gridded precipitation dataset will provide a basis for the 2D-mesoscale downscaling (WP2.3), it allows for the evaluation of regional reanalyses (WP2.4) and it will be exploited for the identification of significant climate anomalies in recent decades, including the occurrence of heavy precipitation events (WP3).

**DWD** will make the European window of the GPCP dataset available for the EURO4M products and services. The GPCP provides global analysis of precipitation on the earth's land surface based on *in situ* rain gauge data. This is particularly relevant in order to place the European results in the global context.

**KNMI** will extend and update the ENSEMBLES gridded daily dataset for Europe (E-Obs) at a resolution of 25 by 25 km (Haylock *et al.*, 2008). This dataset is developed with participation from the EUMETNET-ECSN community (see Section 2.3). It is primarily based on the station data from the ECA&D project and the ENSEMBLES dataset is now maintained by KNMI. The E-Obs dataset is recognized as Europe's baseline dataset to serve a (official) WMO –RAVI Regional Climate Centre functionality for high resolution observational data. Efforts will be made to improve the spatial coverage of the daily dataset and include more variables (in particular variables required for downscaling in WP2, such as snow cover) plus better metadata (partly using Google Earth and GIS). Methods for homogenizing daily time series will be tested and applied, building upon the flagging technique which is currently in use in ECA&D. This activity is linked to the COST-Action HOME: Advances in homogenization methods of climate series: an integrated approach, which is led by MF (see [www.homogenisation.org](http://www.homogenisation.org)). The gridded E-Obs dataset which covers all of Europe will be updated at 6-monthly time intervals during the course of the project, but near-real time in case of emerging extreme event.

**MF** will provide some of its own regional datasets in support to the project and apply its own quality control procedures to the collected data. Notably, MF will extend the different controls such as thresholds, inter parameter consistency, time evolution and spatial consistency to the collected data. Additionally, MF will try to improve such controls by using, first over France and then when possible over other regions of interest, additional data such as radars, models and satellites in order to qualify in the best way ground measurements used by the project (notably for WP2).



**UEA** will use the latest version of the CRU monthly high-resolution ( $0.5^\circ$  by  $0.5^\circ$  latitude/longitude) datasets (CRU TS 3, an update of CRU TS 2.1, Mitchell and Jones, 2005) to assess the quality and accuracy of many of the gridded dataset products (including those based on ERA-40) available for Europe. In the course of development of the CRU high-resolution datasets (which include the following variables: maximum and minimum temperature, precipitation, rainy day counts, vapour pressure and cloudiness), UEA has gained considerable experience in the “best practice” of gridding and interpolation of station data. For example, it is rarely optimum to use the data expressed in the original measured units. Additionally, many users take CRU products to derive other fields, but the approaches employed are generally sub-optimal. With these aspects in mind, UEA will develop two additional fields from CRU data products – Potential Evapotranspiration (PET) and Palmer Drought Severity Index (PDSI) fields, both of which will be useful for regional climate model validation, but more importantly for the online monitoring of drought across Europe (WP3).

**MO** will make a link between this WP and the data work MO is already doing as part of ongoing projects (including the planned global reanalyses) and as part of the international ACRE initiative. In particular, this work will include improvements to HadISST forcing fields and sub-daily SYNOP data in collaboration with NCDC.

#### *WP1.2: Developing gridded datasets based on long-term remote sensing data (DWD)*

The use of satellite-derived data for monitoring the climate system is a relatively new application. The SAF on Climate Monitoring (CM-SAF), which is under the guidance of DWD, is fully dedicated to the high-quality long-term monitoring of the climate systems state and variability. However, the overall capacity of satellite-derived data for climate monitoring has not been evaluated and used satisfactorily. EURO4M will enable the use of satellite data for climate monitoring by constraining their uncertainty with ground truth (*in situ* observations). EURO4M therefore adds value to already existing products of EUMETSAT-SAFs in order to build a consistent and continuous monitoring capability for ECVs.

**DWD** will evaluate the capacity of the SAF products and methods for climate monitoring and will add satellite-based gridded datasets to the regional observation system. DWD will provide 20 yr of water vapour data, cloud albedo, surface solar radiation and surface albedo. These datasets will be used for the evaluation of the regional reanalysis in WP2, but also for multi-purpose products and services in WP3. Surface solar radiation, surface and cloud albedo will be based on the well established Heliosat method using MVIRI images (Beyer *et al.* 1996; Hammer *et al.*, 2003; Drews *et al.*, 2008). From 2005 onwards, the Heliosat based surface solar radiation will be merged with the MS and KNMI dataset in order to take benefit of the improved capabilities in monitoring of clouds from the SEVIRI satellite instrument on-board of MSG. The MS dataset is based on an optimized Heliosat method applied to SEVIRI for the Alpine region, characterised by an improved cloud and snow discrimination and consideration of topography effects (Dürr and Zelenka, 2008; Dürr *et al.*, 2008). Thermal radiation fluxes at the earth surface will be used to supplement the satellite-based shortwave radiation fluxes in order to generate a long time series of surface radiation budget.

The GPCC precipitation dataset (Rudolf and Schneider, 2005, Beck *et al.*, 2005) is a well established global dataset of precipitation based on ground measurements. However, it lacks information over the ocean. This drawback can be overcome by data fusion of the GPCC dataset with the HOAPS precipitation retrieved from satellites over the ocean. The Hamburgs Ocean Atmosphere Parameters from Satellite (HOAPS) has been developed at the MPI Hamburg (Grassl *et al.*, 2000). The algorithm to retrieve precipitation is presented in Bauer and Schuessel (1993). It will be investigated if the spatial extrapolation of *in situ* precipitation measurements (GPCC) can be improved by the use of the KNMI satellite derived precipitation dataset provided in WP 1.2, especially in regions with relative low density of ground measurements. A method for appropriate data fusion of these datasets will be developed and applied in order to provide the user community a optimal global precipitation dataset which covers at least 20 years.

Finally, DWD will provide a 4.5 y dataset of water vapour based on the International ATOVS Processing Package (IAPP, Li *et al.*, 2000) and a 20 yr water vapour dataset over the oceans based on HOAPS.

**KNMI** will deliver high-resolution datasets of cloud properties, precipitation and surface solar irradiance (SSI). Cloud properties (optical thickness, particle effective radius, and water path) are retrieved routinely within the CM-SAF from MSG-SEVIRI observations using the Cloud Physical Properties (CPP) algorithm (Roebeling *et al.* 2006). SSI and precipitation constitute the most important effects of clouds at the surface. Their observation from space complements the surface measurement records because of the high spatial resolution and better coverage of satellites. Precipitation occurrence and intensity are derived from retrieved cloud properties, showing good agreement with weather radar observations over the Netherlands (Roebeling and Holleman, 2008). SSI is determined by combining the cloud property retrievals with other variables, most importantly total precipitable water. The SSI product has been validated with pyranometer measurements in the Netherlands (Deneke *et al.*, 2008). Data will be delivered for the MSG time frame (2005 to present), which may be extended backwards in time in a later stage. Additional validation activities for SSI and precipitation will be performed to assess the quality of the products outside the Netherlands. Applications of the datasets include verification of reanalysis data (WP 2), downscaling (WP 2) and guidance for spatial interpolation of in-situ measurements (WP 1).

**MO** will generate maximum and minimum surface air temperature products from satellite data available in the Land-SAF. These will be made available along side the gridded air temperature datasets based on surface observations (see ECV: Surface Temperature in Section 1.2). These efforts will result in a blended satellite/in situ dataset, which will aid heatwave analysis, etc. In addition, sunshine hours will be estimated. The results will be used in WP 3.

WP1.3: *Data coordination and access to national archives (URV)*

Although past and ongoing European and national funded projects, together with international and national bodies' initiatives, have increased the availability of historical climate data over Europe (mainly for air pressure, temperature and precipitation data), the current availability of historical surface climate data is still highly inadequate. This is particularly true over key but scarcely documented regions, such as southern and eastern Mediterranean, as well as for other ECVs than temperature or precipitation. Therefore, it is still urgently needed to fill the gap for surface climatological data and enhance the availability of long-term/high-quality surface climate data at the finer temporal and spatial scales.

While some of the existing components of the atmospheric observation system are truly operational, several of the ground-based *in situ* and remote sensing networks are funded and operated by agencies in support of research programmes; so their continuing operation is by no means assured. Similarly, many of the data analysis and assimilation systems are developed and funded on research budgets. Often, the data from these systems are not made available and included in integrated datasets (they remain stand alone). The assessments of their (potential) contribution to the European atmospheric observation system will be pursued in this WP, including institutional and international aspects.

**URV** will work together with GMES-ISOWG, GCOS, ECSN, ACRE, MEDARE (which URV lead), etc., in order to identify deficiencies of the European long-term climate data availability and accessibility, coordinate new actions for targeted data rescue over Europe and ensure an enhanced accessibility to climate data produced by past and ongoing European and international projects and national archives. URV will carry out a cost/benefit analysis of undertaking Data Rescue (DARE) activities and assess availability/accessibility of historical climate datasets for relevant ECVs. Specifically, it will work together with the MEDARE community to improve accessibility to historical climate data over the Greater Mediterranean Region (GMR), in general, and over eastern and southern Mediterranean arid and semi-arid areas, in particular. Finally, URV will subject the new recovered/digitized ECVs records to a quality control and homogenization assessment (using the techniques recommended by the COST-Action HOME) in order to ensure they are of high-quality. The data which are made available will be included in the gridded datasets of WP 1.1.

**MO** will contribute to this effort and use as its underpinning framework, existing linkages to the international ACRE initiative plus specific projects covering parts of the European region - EuroCryoClim in the Arctic, EUMETNET-ECSN for Europe, and MEDARE, MedCLIVAR and the FP7 CIRCE project for the Mediterranean. It will also link to other regional to sub-regional projects such as BALTEX (including the Baltic Assessment of Climate Change), Nordic climate co-operation (NORDKLIM), and The International Association for the Promotion of Co-operation with Scientists from the New Independent States of the Former Soviet Union (INTAS), especially the project: Snow Cover Changes over Northern Eurasia during the last century (SCCONE). **MO** will work closely with international data rescue and recovery initiatives such as WMO DARE and similar activities within CLIVAR and GCOS (e.g. the GCOS Regional Action Plan for the Mediterranean Basin).

**NMA-RO** will liaise with NMHSs in Eastern Europe to obtain better access to the national climatological archives in this sub-region. Earlier experience has shown that assigning national database experts only a simple data provider role is counterproductive. The national experts know the regional characteristics and limitations of their archived data and this knowledge can add value when integrating these data into European datasets. The correct treatment of access to the national archives will form an example of good practice in the sensitive domain of data exchange, demonstrating a win-win situation for all stakeholders. Also for capacity building, a more active involvement of national climatologists is crucial. NMA-RO will facilitate integration of additional data in European datasets.

### **Deliverables**

- gridded high-resolution daily precipitation dataset for the Alpine region and analysis of daily to decadal precipitation variations (D1.1, M1.1, D1.2);
- European window of the GPCP dataset available for the EURO4M (D1.3);
- extended and updated ENSEMBLES gridded daily dataset for Europe (D1.4, M1.2, D1.5);
- new UEA/CRU data products for Potential Evapotranspiration (PET) and PDSI (D1.6);
- new satellite-based gridded datasets based on CM-SAF (D1.7, M1.3, D1.8), MSG (D1.9, D1.10), Land-SAF (D1.11);
- datasets for the Mediterranean (D1.12, D1.13, M1.4);
- proposal for additional data rescue activities required (D1.14).

Table 1.3d Work package description – part 2

Work Package number	WP2	Start date or starting event:					Month 1
Work Package title	Regional reanalysis						
Activity Type							
Participant number	1	2	3	4	5	6	7
Participant short name	KNMI	MO	URV	NMA-RO	MS	DWD	SMHI
Person-months per participant		111			18	27	41
Participant number	8	9					
Participant short name	UEA	MF					
Person-months per participant	20	64					
<p><b>Objectives</b></p> <ul style="list-style-type: none"> <li>• develop a state-of-the-art system capable of a comprehensive regional reanalysis and 2D downscaling;</li> <li>• deliver a multi-year dynamical reanalysis for Europe with a less advanced system;</li> <li>• evaluate the newly developed system and the multi-year dataset produced with the less advanced system using independent observational datasets;</li> <li>• review available observations and prepare for a longer reanalysis period</li> <li>• improve capability for reanalysis through better input data</li> </ul> <p><b>Sub-WPs</b> (with lead participant in brackets)</p> <p>WP2.1: <i>Building capability for advanced regional data assimilation (MO)</i></p> <p>WP2.2: <i>Dynamical downscaling of ERA (SMHI)</i></p> <p>WP2.3: <i>2D-mesoscale downscaling (MF)</i></p> <p>WP2.4: <i>Evaluation (MS)</i></p> <p>WP2.5: <i>Improvement of input data for reanalyses (UEA)</i></p>							

## Description of work

### WP2.1: *Building capability for advanced regional data assimilation (MO)*

Although a multi-decade regional reanalysis with a state-of-the-art assimilation system will be made as part of the project (see 2.2), this work package will make significant further improvement towards setting a very advanced regional reanalysis system in place and demonstrate its capabilities.

**MO** will develop a regional atmospheric data assimilation reanalysis system, and demonstrate its effectiveness. Thanks to the Unified Model system, the model used will be state-of-the-art for both climate and NWP applications. The system will be based on the Met Office Hadley Centre's regional climate model and the regional assimilation used for its operational North Atlantic and Europe (NAE) NWP configuration. The September 2008 operational NAE assimilation uses 4D-Var with a 12 km/36 km resolution, plus latent heat nudging of precipitation data. R&D is underway for an upgrade in 2009, which will replace the latent heat nudging by 4D-Var assimilation of the same surface precipitation estimates. The observational database used will be that gathered at ECMWF for ERA, enhanced by other packages in this proposal, in particular 2-dimensional accumulated precipitation estimates. Boundary conditions will be from ERA-Interim (or a later reanalysis if available). SST will be that used in ERA-Interim. Since there is significant development effort in this activity, the results will not be in time for feeding into further downscaling in WP2.3.

### WP2.2: *Dynamical downscaling of ERA (SMHI)*

A current state-of-the-art dynamical downscaling for an extended multi-year period will be made as part of EURO4M with a comparatively cheaper system computingwise, than developed in WP2.1. In this way, the first results of regional reanalysis can already be used within EURO4M for the products and services delivered in WP3.

**SMHI** will complement the dynamical reanalysis by MO using the HIRLAM system. SMHI will perform HIRLAM 25 km dynamical reanalysis (HIRLAM-A programme). It will employ the HIRLAM 3D variational analysis and the HIRLAM grid point model for the assimilation (Gustafsson *et al.*, 2000 and Lindskog *et al.*, 2001). Conventional observations (SYNOP, AIREP and AMDAR, buoys, TEMP) will be used as well as METEOSAT wind observations. The observed data will be analysed in 6 h windows and analyses will be produced every 6 h. The lateral boundary conditions will come from the global ERA-40 (1960-1988) or ERA-Interim (1989-2010) reanalysis. A large scale forcing may be applied every 12 or 24 h from those datasets, mainly in order to benefit from the use of satellite radiances in the ERA reanalyses which involved a lot of quality control and bias corrections. Those data are important for defining the large scale features present also inside the Limited Area chosen. This large scale forcing can be replaced by a large scale constraint,  $J_k$  that is being developed in HIRLAM. The HIRLAM reanalysis provides a downscaling from global resolution (125 km for ERA-40 and 80 km for ERA-Interim) to 25 km in the Limited Area. The increased resolution provides more detail and realism due to both the 3D-VAR analysis of observations but also to a great deal from the 25 km grid point model used in the forecast steps of the assimilation.

### WP2.3: 2D-mesoscale downscaling (MF)

The 2D-mesoscale downscaling techniques used in this WP provide added value for surface and near-surface fields, because they are able to include additional observations and work in higher resolution than the regional reanalysis.

**SMHI** will develop a 2D-downscaling system where features from the present SMHI system MESAN are utilized together with an improved description of predictors in combination with developments within the SAFRAN system at MF. In a pilot project in 2007, SMHI applied the present MESAN system to a 2D-downscaling for Europe (ERAMESAN) using ERA-40 as background (first guess) fields. In this WP the results from WP2.2 will be used as background fields for the 2D-downscaling. Possible spatial resolution for the resulting dataset would be of the order of 3-12 km depending on variable. An improved MESAN (also used in ERAMESAN; Jansson *et al.*, 2007) meso-scale analysis system will use many more observations than those used in the 3D-VAR. The advantage of the MESAN is that it is 2-Dimensional (only) and geared specifically for a faithful gridded analysis of surface and near surface fields for nowcasting and climate analysis. The present structure functions, which have been developed with basic parameterizations for topography and water cover, will be further elaborated through scaling for predictors like topography, distance to sea etc. The necessary first guess fields will be obtained from the dataset that results from WP2.2. Observations to be used are data from ECMWF plus other quality controlled data which will be made available in a usable format by WP1. Products will comprise not only the model-assisted analyses of ECVs produced by data assimilation, but also metadata providing information on the quality control and bias-correction of the assimilated satellite and *in situ* data. These will be used in WP3 for uncertainty information.

**MF** will downscale the 3D reanalysis from WP2.2 using low-level datasets, 2D analysis techniques, and land surface modelling. This will provide added value through additional observations and over orography. MF will also improve the 2D downscaling software, in particular the data QC and interface with the 3D reanalysis. As the next step, depending on the results of evaluations in WP2.4, MF in collaboration with **SMHI** will incorporate either some developments (the hourly distribution of precipitation) or specific ideas for mountain areas from the SAFRAN software in the MESAN system. Some developments already used in some NWP systems such as: new structure functions to improve the coastal and the topography effects (Navascuès, 1997; Rodriguez *et al.*, 2003), snow analysis using the classical observations and satellite pictures will be investigated. The new system MESAN-SAFRAN will be able to also use the long time series of observations that are not used in NWP, including minimum and maximum temperature, etc. The datasets for this activity will come from WP1. Finally, the new system will be coupled with a surface and hydrological scheme in order to improved surface variables and in particular the climatology of snow cover.

#### WP2.4: Evaluation (MS)

The activities in this WP aim at evaluation of the different reanalysis/downscaled datasets. It brings together the evaluation activities of most of the partners involved in the production of the gridded datasets. The station-based and satellite-based datasets developed in WP 1 will be used together with existing other datasets and reanalysis products. The goal is to improve the reanalysis downscaling systems and to provide uncertainty information for the datasets that are used in WP 3. One issue that will receive attention when synthesising the results of this WP is the fact that some of the downscaling systems have been calibrated and tested for one particular region or country. How do they perform elsewhere in Europe?

**MS** will evaluate the regional reanalysis with respect to the representation of daily to decadal precipitation variations in the Alpine region. As a reference, the high-resolution gridded precipitation dataset developed in WP 1.1 will be used. The high station density underlying this dataset will permit for an accurate upscaling to the resolution of the regional reanalysis, which is essential when evaluating characteristics of heavy precipitation. Special consideration will also be given to the uncertainties of the gridding (spatialisation) procedure when assessing the reanalysis.

**MO** will demonstrate the value of the system developed in WP 2.1 by comparing a year or two of results from: ERA-Interim; the Met Office regional climate model nested in ERA-Interim; assimilation of ERA-Interim observations into the same regional model; assimilation of ERA-Interim observations and processed precipitation observations; downscaling of ERA-Interim and assimilation of some observations in a different model in WP 2.2.

**SMHI** will compare the NAE-based reanalysis system under development at the MO (WP 2.1) against the state-of-the-art HIRLAM-based system used for the multi-decade reanalysis at SMHI from (WP 2.2). This comparison will be important to assess the possible benefit from a more advanced system for future regional reanalysis. The comparison is possible towards the end of the project. This leaves open the future possibility of a multi-model reanalysis as a means of assessing uncertainty.

**MF** will build on earlier evaluation of ERAMESAN (Jansson *et al.*, 2007) by SMHI, which focussed on the territory of Sweden where most observational data were available. MF will concentrate on another area and compare ERAMESAN with SIM (SAFRAN-ISBA-MODCOU; Habets *et al.*, 2008; Qunitana-Segui *et al.*, 2008) analysis over France for the time-period 1980-2004. A more local comparison in a mountainous area will be done with the SAFRAN-NIVO analysis, which is considered as a “reference” over the Alps (Durand *et al.*, 2008). This will provide inside of the gain of more observational data and a higher resolution system. More importantly, the newly developed MESAN-SAFRAN system will be evaluated. The snow albedo, computed by the snow scheme, will be compared with Land-SAF data (WP 2). A comparison of the surface fluxes (latent and sensible heat) with in-situ measurement available in Europe (Cabauw, Sodankyla, Lindenberg) will demonstrate the “physical” realism of the system.

**DWD** will verify the reanalysis data for cloud parameters, surface solar radiation, surface albedo, water vapour, precipitation and top of atmosphere radiation flux densities by comparing to existing CM-SAF products and products generated within Wp 1.2. An example of verification of reanalysis data is published in Babst *et al.* (2008). DWD will support satellite inter-calibration activities in order to support the generation of homogenized and inter-calibrated radiances for specific satellite sensors. DWD will investigate the effect of satellite calibration uncertainties and inhomogeneities on cloud, radiation, temperature and humidity reanalysis products (including precipitation). The results will be analysed and suggestions for improvements will be drawn and reported to the reanalysis centres.



### WP2.5: Improvement of input data for reanalyses (UEA)

ERA-40 (and NCEP/NCAR) use as input files historic satellite and *in situ* (radiosonde and surface) data. These files are essentially archives of all the operational input to weather forecasts over many decades and have been put together by NCEP/NCAR. No changes have been made to these operational inputs since the data were collected. Operational analyses reject obvious outliers in real time, but over the last 50 years there have been considerable advances in the homogeneity testing of climate data (see e.g. Thorne *et al.*, 2005a,b and Haimberger *et al.*, 2008). The purpose of this WP is to a) determine exactly what is in this operational archive (it has only ever been used as reanalysis input data) and b) to replace and upgrade, where this is possible, these data with improved datasets available nationally and internationally. If we expect subsequent generations of reanalyses to be improvements on earlier versions, it seems essential that we make the effort to improve the basic input these expensive reprocessing operations require. As climate models underlying each reanalysis improve with time, then so should the basic input data.

UEA together with MO (and in cooperation with the institutes performing the reanalysis) will analyse the improvement of capability for reanalyses through better basic input data. A major example of this is the important radiosonde input. A number of papers (*op. cit.*) have made significant advances in adjusting radiosonde data for biases resulting from changes to sensors, instruments etc – adjustments which can be in the range 1-5°C for temperature and larger in relative magnitude for humidity (greater at higher elevations).

There is also a dramatic increase over time in the amount of data available for assimilation in reanalysis, especially after the introduction of operational satellite observing systems in the 1970s. The data assimilated in ERA-40 will be reviewed, and their importance, deficiencies and any remedial action already undertaken since ERA-40 will be tabulated for different epochs. Improvements in historic satellite datasets are expected to result from data reassessments (e.g. against 20<sup>th</sup> Century Reanalysis Project results) and reprocessing by EUMETSAT and other satellite agencies (see also WP 1.2). Emphasis in this WP will accordingly be placed on *in situ* data, in particular for the pre-satellite period. The sets of surface and radiosonde data assimilated in ERA-40 will be assessed with regard to the amounts of data they contain, the spatial coverage and the periods of time covered. This will guide which of the many other archives of synoptic surface and radiosonde data should be evaluated, and if necessary reformatted or digitized, to improve the quantity and quality of conventional data for assimilation in future reanalyses. The principal surface archive is ISH (see earlier under the ECV – surface humidity) and we will assess the degree of commonality between this and the surface component of the reanalysis archive. It is also expected that the HadAT radiosonde archive (Thorne *et al.*, 2005a,b) will prove useful for homogeneity assessment to minimise time-dependent biases with sondes, together with additional information from reanalysis feedback files (Haimberger, 2005). Homogeneity will initially consider temperature measures, but will extend to the humidity sensors as progress in other projects permits. The IGRA archive of individual ascents will be accessed to determine how many extra ascents can be added, provided several quality criteria are met. Emphasis in these homogeneity assessments and additional data will be on Europe for the period prior to 1979, but will extend to other land regions of the world where sonde coverage is poor. In particular, the entire British Antarctic Survey sonde archive for the Antarctic will be added. In determining which additional data need to be digitized there will be active consultation with WP 1.

Bias adjustment of much of the basic satellite input data has also made advances since its initial operational use (the MSU series of instruments is a good example here). Many similar adjustments will also have been made to daily surface temperature and pressure data, and improved datasets can be incorporated. Considerable additional data have also been digitised from the 1950s-1980s that were not internationally exchanged at the time, and these will also be added to the data bank.

### **Deliverables**

- new state-of-the-art NAE-based regional atmospheric data assimilation reanalysis system (D2.1, M2.1, D2.2);
- HIRLAM-based reanalysis dataset at 25 km for most of the past 20 yr period (D2.3, M2.2)
- improved MESAN downscaled dataset at 3-12 km resolution for most of the past 20 yr period (D2.4);
- new MESAN-SAFRAN downscaling system (D2.5, M2.3, D2.6);
- evaluation reports for specific components of the different (downscaled) datasets and systems (D2.7, D2.8, D2.9, D2.10, D2.11, D2.12);
- synthesis report for all evaluation activities in EURO4M (D2.13, M2.4);
- analysis of the improvement of capability for reanalyses through better basic input data (D2.14);
- access to in situ datasets not used in existing reanalysis (D2.15).

Table 1.3d Work package description – part 3

Work Package number	WP <sub>3</sub>		Start date or starting event:				Month 1
Work Package title	User-oriented information and climate change products						
Activity Type							
Participant number	1	2	3	4	5	6	7
Participant short name	KNMI	MO	URV	NMA-RO	MS	DWD	SMHI
Person-months per participant	31	12		36		28	
Participant number	8	9					
Participant short name	UEA	MF					
Person-months per participant							
<p><b>Objectives</b></p> <ul style="list-style-type: none"> <li>• produce innovative and integrated climate change products that are multi-purpose, but based on user requirements, and form the basis for future GMES implementation services</li> <li>• develop regular, flexible and holistic reference baseline CIBs (see Section 1.1) that are generic in the sense that they serve the full range of climate users and applications sectors in Europe within the wider global community;</li> <li>• manage and operate the European Climate Liaison Team (CLT) and implement a user feedback loop by compiling user requirements for all GEO areas (e.g. define the most crucial climatological parameters and how they should be summarized) and providing scientific guidance on the best ways to disseminate and use the EURO4M products and services;</li> <li>• link to the WMO Information System (WIS), which will form one operational exemplar of the GEOSS architecture implementation process;</li> <li>• support the user community with active training, e-learning/training modules;</li> <li>• work on capacity building closely with candidate countries and developing countries, which will be among the largest potential beneficiaries of international co-operation in atmospheric observation.</li> </ul> <p><b>Sub-WPs</b> (with lead participant in brackets)</p> <p>WP<sub>3.1</sub>: <i>Climate Indicator Bulletins</i> (KNMI)</p> <p>WP<sub>3.2</sub>: <i>Climate Liaison Team</i> (DWD)</p>							

## Description of work

### WP3.1: *Climate Indicator Bulletins (KNMI)*

Climate Indicator Bulletins (CIBs) are examples of sound user-oriented multi-purpose products for future GMES services using the output from WP1 & WP2. These CIBs will provide scientific input for policy implementation and near-real time reporting for emerging extreme events. In consultation with key user groups via WP3.2, this WP will develop integrated data products and services based on ECVs for the region of Europe, and where appropriate the Northern Hemisphere and globally. Among the products are climate indicators at the appropriate level of aggregation and processing for the main user applications. Clearly, these CIBs must be scientific, but importantly they must also be easy to understand – in relation to a question/decision/activity. After the end date of the project these can all become part of the GMES implementation services on climate change.

**KNMI** together with **NMA-RO** and **MO** will produce the CIBs and identify and summarise, in a non-technical manner, significant climate anomalies and features including uncertainty estimates. Dramatic examples which will be used to illustrate potential CIB impact are the summer of 2002 with widespread flooding in central and eastern Europe, the European heat wave of 2003, the European summer of 2005 with wet and dry extremes, the winter of 2005/2006 with cold and warm extremes, the winter of 2006/2007 as the very likely warmest for more than half a millennium, and the summer of 2007 with widespread drought and fires in large parts of the central and eastern Mediterranean. This WP will revisit these events, and describe how the use of monitoring information will significantly improve with this project in place.

Experience shows, that after an extreme event (winter storm, flooding) in near-real time there is usually only information available from individual countries. The larger scale picture is missing and cannot simply be derived from raw data. Thus an advantage of the baseline CIBs would be to provide, in near-real time, a larger scale continental picture of weather/climate extremes that complements the national information provided by NMHSs. The same is true for longer lasting extremes, such as heat waves or drought periods. With EURO4M, we aim to be able to quickly and efficiently compile and present a broader, higher detailed picture of European climate anomalies and their impacts than was previously possible.

CIBs will be published both as bulletins (electronic documents/newsletters) as well as being provided through a web-portal, including access to the associated data series, gridded datasets and editorial texts. The information will be based directly on the climate change time series of ECVs made available in WP1 and WP2. Note that the “optimum” information will rarely come consistently from a single source. The system for producing CIBs will be made configurable and flexible, and capable of reacting in near-real time. CIBs will both cover the whole of Europe as well as sub-regions within Europe. Aggregated information will also be provided for individual countries, large river basins and Europe as a whole.

### WP3.2: *Climate Liaison Team (DWD)*

As a potential future GMES service on climate change monitoring, EURO4M must interface directly with the full range of intermediate- and end-user requirements, including disaster prevention, health, energy, water resources, ecosystems, forestry agriculture, transport, tourism and biodiversity. The Climate Liaison Team (CLT) developed in this WP will establish a communication process, which responds to information flow in both directions and evolves through time, so that users can both obtain and influence the nature (user driven data format, content and delivery style) of the information they need while understanding the strengths and limitations of the monitoring products. In doing so, the team will establish sustainable links between EURO4M and all GMES services and GEO societal benefit areas.

**DWD** will co-ordinate the CLT as a cross-cutting activity of the project. Through their mediation, the user-oriented results of EURO4M research will feedback directly into applications and impact assessments relevant to European societal and community needs. The stakeholders (impact community and decision makers) needs in terms of climate data and products cover a large range of raw or derived variables, indicators, spatial and temporal scales, uncertainty assessment at various levels of complexity. The stakeholder community is however diverse and not well organized. This will make it difficult to obtain uniform feedback for the multi-purpose products. Therefore DWD will review existing reports and information about user needs and will generate a meta user-requirement document, specifically addressing the intersection of user needs across the range of user communities. DWD will also generate a data storage and dissemination document, describing the user driven and INSPIRE-compliant infrastructure used for these tasks in WP1 and WP2 (see Section 2.4). This will provide necessary links to the WMO Information System (WIS), which will form one operational exemplar of the GEOSS architecture implementation process. Information gaps will be closed by generation and evaluation of specific surveys. Actors in the relevant core and downstream GMES services and in related national and international programmes will be actively involved in this process. The lessons learned from the process for establishing user needs that has been used for the selection of fast track and pilot services will be considered too.

**DWD** will demonstrate the usefulness of EURO4M services to policy-makers, researchers, planners and the citizens and their organizations. Among others, video podcasts will be generated for the public and environmental assessment reports covering different skill levels. Training material will be developed in order to promote the benefit of integrated use of climate data. A course management system (MOODLE), designed using sound pedagogical principles, will be used to create effective online learning modules. The CM-SAF group at DWD has already established training facilities for operation in an international context. For example, CM-SAF organises in cooperation with EUMETSAT a series of international training workshops on the use of satellite data for climate applications.

#### **Deliverables**

- Climate Indicator Bulletins (CIBs) as sound user-oriented multi-purpose products for future GMES services that synthesize all existing information relevant for environmental assessments and respond quickly (within a few days) to climatic events (D3.1, D3.2, M3.1);
- Climate Liaison Team (CLT)(D3.3, M3.2);
- a pan-European meta user-requirement document for the intersection of user needs across the range of user communities and a data storage and dissemination document, describing the user driven and INSPIRE-compliant infrastructure of EURO4M (D3.4, D3.5);
- video podcasts, training material and workshops on monitoring of climate extremes, also in developing countries (D3.6, D3.7, D3.8).

Table 1.3d Work package description – part 4

Work Package number	WP4	Start date or starting event:					Month 1
Work Package title	Project management, coordination and sustainability						
Activity Type							
Participant number	1	2	3	4	5	6	7
Participant short name	KNMI	MO	URV	NMA-RO	MS	DWD	SMHI
Person-months per participant	12	3		4			
Participant number	8	9					
Participant short name	UEA	MF					
Person-months per participant	3						
<p><b>Objectives</b></p> <ul style="list-style-type: none"> <li>• coordinate the consortium and all EURO4M activities; manage day to day operation of the project; ensure that deliverables are completed on time;</li> <li>• ensure that the EURO4M atmospheric observation system is maintained and operated for the duration of the project, systematically generating timely ECVs and CIBs;</li> <li>• prepare realistic plans to sustain and maintain the EURO4M generic reference baseline CIBs as a viable entity beyond the lifetime of the project and as a future operational GMES service on climate change monitoring.</li> </ul>							

### **Description of work**

The management structure has been designed to match the complexity of the project and manage effectively the level of integration required through the project's lifecycle. The management of EURO4M will be an effective structure intended to maximize the contribution to the European Environment Earth Observation System. It includes: the Coordinator **KNMI** responsible for overall coordination of the project, and reports to the European Commission; the Management Board with leaders of each WP responsible for the overall management of the project. The general assembly consisting of one representative from each participant institution will convene annually to discuss EURO4M progress and plans. Other stakeholders from the wider network and representatives of related research programmes will be invited to attend the general assembly and contribute to the planning process.

A coherent management structure delivering effective supervision of the project on a day to day basis is essential if the various strands of the project are to be properly combined, and the exciting potential for new knowledge is to be fully realised. Accordingly, management activities have also been given their own WP within the structure of the project as described in Section 2.1. This WP has the responsibility for the overall coordination of the EURO4M project. Its primary objective is to ensure that the activities carried out in the other WPs under the leadership of **UEA, MO** and **NMA-RO** are fully integrated towards a common purpose and hence that the project overall delivers the full benefits to the users and wider community of the extensive and innovative research and training carried out in the project. WP4 will also report project progress to the Commission.

In the final stage of the project, a plan will be set up to define what it takes to secure the availability of the developed GMES services and of the critical observational data beyond the lifetime of the project. In this way a smooth transition from the demonstration stage with R&D funding to a pre-operational and operational stage will be possible.

### **Deliverables**

- regular MB meetings and General Assemblies (D4.1, D4.2, D4.3, M4.1);
- EURO4M website, brochure and logo (D4.4, D4.5);
- activity reports and management reports for the Commission (D4.6);
- detailed implementation plans (D4.7);
- plans to sustain and maintain the EURO4M generic reference baseline CIBs as a future operational GMES service (D4.8);
- major final EURO4M conference (D4.9).

**Table 1.3e Summary of staff effort**

Participant no./short name	WP1	WP2	WP3	WP4	Total
1. KNMI	32		31	12	75
2. MO	24	111	12	3	150
3. URV	42				42
4. NMA-RO	36		36	4	76
5. MS	36	18			54
6. DWD	16	27	28		71
7. SMHI		41			41
8. UEA	31	20		3	54
9. MF	18	64			82
<b>Total</b>	<b>235</b>	<b>281</b>	<b>107</b>	<b>22</b>	<b>645</b>

### ***Graphical presentation of the components***

As illustrated in Figure 1.3c, the activities in the 4 WPs of EURO4M strongly interact. In order to enable EURO4M to accomplish its goal of developing the capacity for, and delivering the best possible and most complete (gridded) climate change time series and monitoring services for Europe a total of 4 WPs is required. WP 1, 2 and 3 form the face of the project and fit together tightly. WP 4 is overarching the project ensuring adequate project management (see Section 2.1).

**In summary**, WP 1 and WP 2 produce the data that will feed into WP 3. Many of the WP 1 observations will also feed into the reanalysis activities in WP 2. WP 3 will produce the EURO4M multi-purpose products and services. For these to be state-of-the-art, we need improved regional observation datasets (WP 1), as well as advanced regional data assimilation and reanalyses (WP 2). WP 3 also has prime responsibility for project outreach.

**In detail**, WP 1.1 will provide high-quality gridded ground truth observations feeding the CIBs (WP 3.1) and serving as input for the evaluation of the regional reanalyses (WP 2.4). Moreover, these datasets constitute an indispensable basis for the calibration and application of 2D-mesoscale downscaling in WP 2.3 and satellite retrieval algorithms in WP 1.2. The satellite-based datasets of WP 1.2 also feed into the CIBs and will be used for reanalysis evaluation too. The data coordination activities in WP 1.3 make additional *in situ* datasets accessible to WP 1.1 and interact with the *in situ* component of GMES.

WP 2.1 guarantees that the possibilities of regional reanalysis as an optimal integration tool for ECVs are fully exploited and links to global reanalysis outside the project. The downscaling activities in this WP and in WP 2.2 and 2.3 are jointly evaluated in WP 2.4. This evaluation is partly based on the datasets developed in WP 1.1 and WP 1.2. It enables a rigorous quantification of the capabilities of regional reanalysis and an assessment of the various methodological options in downscaling techniques developed and implemented in WP 2.1, WP 2.2 and WP 2.3. The results provide elementary guidance for the appropriate derivation and interpretation of the derived products in the Climate



Indicator Bulletins (WP 3.1). WP 2.5 will provide additional capability for the work in WP 2.1 and WP 2.2 through improved input data for regional (and global) reanalysis.

The dataset production and evaluation activities in WP 1 and WP 2 complement each other. They are brought together in WP 3.1, where the best possible multi-purpose products and services will be developed. WP 3.1 makes the high quality spatial ECV data accessible at the required level of aggregation. In particular, the “optimum” results need to be based on different underlying ECV datasets. The added value given to the raw data by this WP should not be underestimated. The EURO4M integrated climate change products and services will be produced in the form of CIBs. Data products and services in the CIBs will evolve in response to feedback from the various users through user interaction in WP 3.2. The Climate Liaison Team of WP 3.2 will link user needs to the CIBs. The capabilities for meeting specific user requirements will ultimately depend on the quality of the raw data and hence a vivid exchange of information between WP 1, WP 2 and the CLT of WP 3.2 is decisive. In doing this, WP 3.2 will demonstrate that bridging the gap between the climate community and the user community is feasible.




Figure 1.3c. Interactions between the 4 WPs of EURO4M plus links to the outside world (see Section 2.3).

### ***Significant risks and associated contingency plans***

For the successful implementation of the EURO4M aims we rely partly on the cooperation and responsiveness of data holding institutions which are not participating. Overall, we are confident that we can achieve our objectives, as the interest in this project from the community of national responsible institutions and their organisations is very

high. **The links to third party participants already established (in particular EEA, ECMWF and EUMETNET; see Section 2.3) will help to solve any potential risk.**

One of these risks is access to weather and climate data, which is often restricted due to intellectual property rights (see Data exchange). This also affects the results of earlier EU research projects. As a result, major costs may arise from access to meteorological data. The fact that the consortium as a whole represents key climate change data holding institutions in Europe and the fact that the consortium members have been active in most of the related EU research projects in the past places us in the best possible position to overcome this problem. In case data policy restrictions hamper the work, the Coordinator will bring the issue to the responsible parties, including the EUMETNET Council.

In the case that internal disputes will occur during the course of the project, we will begin at the source of the dispute and expand outwards until an equitable solution has been reached. The participant tries to settle the dispute with its respective WP-leader. If not successful, the Management Board (MB; see Section 2.1) will develop a strategy and intervene to settle the dispute with the possibility of sanctions. In case of disagreement within the MB, the Coordinator will take the final decision and he will be responsible for its execution. The Coordinator will report the case and its solution or consequences to the EC appointed project officer.

## 2. Implementation

### 2.1 Management structure and procedures

#### *Management structure*

We propose a layered management structure for the project. This will facilitate the production of deliverables with a minimum of bureaucracy and of business meetings. The structure is designed to ensure that the work of the WPs is carried out efficiently and according to plan, and that this work is then integrated effectively to achieve the goals of the project. **The management structure combines the proven concepts of a number of ongoing and successful EU projects.** With this structure in place, we will be able to react flexibly to unexpected developments which might occur during the lifetime of the project. In addition, a mechanism is planned to open the structure to participants other than the EURO4M core participants.




Figure 2.1a. Management structure of EURO4M.

The following management structure is proposed, as presented schematically in Figure 2.1a:

**General Assembly:** The General Assembly (GA) consists of representatives of all project participants. They will meet on an annual basis. Representatives of the user community, third party participants (see Section 2.3), and Advisory Board (see below) will attend the GA. The General Assembly also forms a platform for direct interaction between the participants in different WPs.

**Management Board:** The top level management of the project will be carried out by the Management Board (MB). The MB will meet at six monthly intervals. It will have responsibility for drawing up the 18 month implementation plan on a rolling basis, review the progress in each WP, and decide on success criteria to proceed with or stop an activity. The MB is also responsible for the coordination of cross-cutting themes. It will be

very important to ensure that these cross-cutting activities are actively coordinated across the WPs in order to avoid duplication and to integrate the results of the project in these areas. The cross-cutting nature of these activities means that they are highly interdisciplinary, involving work in all WPs. This will involve the development of common, clearly-defined unambiguous terminology and definitions of, for example, extreme events and standardization of uncertainty estimates. Managing these cross-cutting activities most effectively will be carried out primarily through small working meetings focussed on these issues and, on a more regular basis, through email discussion groups. Larger workshops on the cross-cutting activities will be organised together with dissemination activities. Part of the EURO4M website will, for example, be dedicated to cross-cutting issues.

The membership of the MB will be made up of all the WP-leaders plus representatives from the EC, GMES, and GEO. The latter will be filled in consultation with the Commission and the GEO secretariat. Currently the WP-leaders are:

**WP1: Regional observation datasets**

Phil Jones, UEA

**WP2: Regional reanalysis**

Andrew Lorenc, MO

**WP3: User-oriented information and climate change products**

Roxana Bojariu, NMA-RO

**WP4: Project management, coordination and sustainability**

Albert Klein Tank, KNMI

**EC Representative:**

To be decided

**GMES Representative:**

To be decided

**GEO Representative:**

To be decided

**Coordinator:** The EURO4M Coordinator leads WP4 (Project Management) and is responsible for overall coordination of the project. He reports to the European Commission and chairs the MB.

**Project Office:** The Project Office will be responsible for the routine administration of the project and the scientific direction. It consists of the Coordinator and support staff at the Coordinators office. The Project Office will monitor progress and facilitate communication between the different groups, organise the scheduled meetings of the project as well as small working sessions between groups when necessary. The Project

Office will also liaise with partners on the financial administration of the project and the provision of necessary documentation to the Commission.

**WP-leaders:** The WP-leaders are responsible for coordinating the various WPs. They also contribute to the top level management of the project through participation in the Management Board.

**Advisory Board:** The Advisory Board will add value to EURO4M through commenting on the overall science plans and directions, and on the progress in achieving the plans. It will consist of representatives from the third party participants linked to the project (see Section 2.3), representatives from collaborative projects (see Section 3.1), as well as representatives from other relevant international agencies and working groups. The Advisory Board will meet once a year and at the start of the project to suggest collaborations. A key task for them is to ensure that the project remains directly relevant to the GMES and GEO process. We will, in consultation with the Commission and the GEO secretariat, fill the positions in this board.

## 2.2 Individual participants

The sections below describe how each of the individual participants is well equipped to deliver her or his required part of the project. Special attention has been given to the responsible scientists involved (bold face).

### **Participant no. 1: Royal Netherlands Meteorological Institute (KNMI), Netherlands**

The Royal Netherlands Meteorological Institute (KNMI) is the national research and information centre for weather, climate and climate change in the Netherlands. KNMI has a long tradition in operational and scientific activities. Climate research at KNMI aims at observing, understanding and predicting changes in the climate system.

**Dr. Albert Klein Tank**, who has been working as a scientist for KNMI for almost 20 years now, will coordinate EURO4M. He is actively involved in observational research embedded in international projects and programmes. Albert Klein Tank leads the EUMETNET project ECA&D ([eca.knmi.nl](http://eca.knmi.nl)) that joins over 40 meteorological services in Europe and the Mediterranean. This project has proved to deliver high quality observational datasets and derived information on indices for studying extremes. He has been involved in the production of the IPCC-WGI-AR4 as a lead author of the chapter on observations. He is currently co-leading the CCI/CLIVAR/JCOMM Expert Team on Climate Change Detection and Indices. In the Netherlands, he is responsible for the national climate change scenario activities at KNMI. His experiences as a project leader and skills to mobilize scientists will be beneficial to managing EURO4M in a successful manner and will ensure an adequate dissemination of the project results. Albert Klein Tank is a member of the FP6 IP ENSEMBLES Management Board, which provides the necessary experience to successfully coordinate EURO4M.

**Dr. Gerard van der Schrier** also is a scientist at KNMI. Following a post-doctoral at the Climatic Research Unit, UEA, where Gerard van der Schrier worked on estimating thermohaline ocean-circulation influences on European climate and on quantification and analysis of European droughts for the 1900-2002 period with Prof. Phil Jones, he returned to KNMI. Working first on climate change from a modelling perspective, by applying a newly developed data assimilation method to model past climate changes, he now focuses on changes in climate extremes from the viewpoint of observations. Gerard van der Schrier's research is largely based on the station data from the ECA&D project for which he also maintains the database and website.

**Dr. Jan Fokke Meirink** holds a PhD in the field of air-sea interaction (2002). From 2001 until 2007 he has been working at KNMI and Utrecht University on the interpretation of satellite measurements of atmospheric trace gases using transport models, and the assimilation of these measurements in order to improve existing estimates of trace gas emissions. The main focus was on methane observations from SCIAMACHY. Since January 2008 he is working in the Climate Observations department at KNMI on retrieval, validation and evaluation of cloud physical properties from MSG-SEVIRI observations. This work is performed in the framework of the CM-SAF.

**Dr. Ir. Rob A. Roebeling** holds a PhD in Environmental Sciences (2008) from Wageningen University, on Cloud Properties Retrievals from Satellite Observations. In 1991 and 1992 he worked at the DLO-Staring Centrum in Wageningen (the Netherlands) on surface flux retrievals from satellite data. In the period 1993-1999 he worked at the Environmental Analysis and Remote Sensing Ltd. as a consultant in the field of boundary layer meteorology and crop growth modelling. He has been project leader of several national and international projects. Rob Roebeling has been employed at KNMI since 2000, where he works as senior scientist on cloud properties retrievals from meteorological satellites. He is principle investigator for the cloud property retrievals within the CM-SAF. Furthermore, he is leading scientific projects in the fields of atmospheric radiative transfer and multi-sensor cloud remote sensing.

## **Participant no. 2: Met Office (MO), United Kingdom**

The Met Office in the UK (MO) is among the world leaders in both NWP and climate research, the latter being grouped in its Hadley Centre. Both use its Unified Model, facilitating projects like EURO<sub>4</sub>M which combine NWP data assimilation with climate studies. The Climate Monitoring and Attribution Group within the Hadley Centre has considerable expertise and experience in the development of high quality historical datasets and gridded products of ECVs for the purposes of climate studies, and their application to observational and model-based studies of climate variations. Several members have been instrumental in leading the development and evolution of key surface datasets that are internationally recognised and utilised (e.g. Central England Temperature, HadISST (Hadley Centre Sea Ice and Sea Surface Temperature dataset), HadCRUT<sub>3</sub> (Hadley Centre and Climatic Research Unit's Air and Marine Temperature Anomalies Version 3 dataset)) for the assessment of climatic variability and change. MO participants have links to the ETCCDI and were also actively involved in the recent EMULATE project, and continue to play a prominent role in ENSEMBLES. The MO Hadley Centre is also a major partner in the international ACRE (Atmospheric Circulation Reconstructions over the Earth) initiative: it is part of the consortium behind ACRE, along with the Queensland Climate Change Centre of Excellence (QCCCE) in Australia, and the National Oceanic and Atmospheric Administration (NOAA) Earth System Research Laboratory (ESRL) and Cooperative Institute for Research in Environmental Sciences (CIRES) at the University of Colorado in the US.

**Dr Rob Allan** is a Climate Scientist at the MO Hadley Centre, where he led the development of surface pressure datasets and products such as HadSLP and EMSLP. Rob Allan is currently the Project Manager of the ACRE initiative and a Co-convenor of the GCOS AOPC/OOPC Working Group on Surface Pressure, with a EURO<sub>4</sub>M third party participant (see Section 2.3) Dr Gil Compo from NOAA/ESRL/CIRES in the US. Dr Compo is leading a component of the ACRE initiative, in which the development of an International Surface Pressure Database (ISPD) is “fuelling” a series of surface-observations-only historical climate quality 3D reanalyses of global weather variables for climate variability and change research. ACRE is helping to facilitate these historical reanalyses through the data rescue of old high-resolution marine and terrestrial observations.

The production of oceanographic ECVs, in particular SST, in near-real time is one of the activities performed within the Ocean Forecasting Research and Development (OFR&D) program located in the MO. The activities of OFR&D also include outreach, with links to the UNESCO-Bilko project and activities such as lessons on the application of remote sensing to oceanography and coastal management. **Dr Nick Rayner** is the Managing Scientist responsible for Marine Data working in the Climate Monitoring and Attribution Group, especially the Hadley Centre Sea Ice and Sea Surface Temperature (HadISST) and the Hadley Centre subsurface ocean analyses datasets.

The members of the Climatic Monitoring and Attribution Group within the **MO** have been involved in the development of UK and/or international climate products and outreach. These include the preparation of quarterly diagrams and information for DECC (UK), monthly graphics and text to update climate information and indices on the MO's Internet site, annual summaries for the UK Royal Meteorological Society's publication *Weather*, contributions to the EUMETNET-ECSN/WMO/DWD Annual Bulletin on the Climate in WMO Region 6, and input to the annual WMO press statement and *Bulletin of the American Meteorological Society* world climate summary.

**Andrew Lorenc** is a Senior Research Fellow of the Met Office. Until recently he headed research and development for NWP data assimilation and ensemble forecasting. He was involved in all three WCRP International Conferences on Reanalysis and was an invited speaker on advanced data assimilation at the last (Tokyo, 2008).

**Dr Bruce Macpherson** manages a team developing operational regional NWP data assimilation systems. He was chair (1999-2004) of working group 3 of the COST Action 717 on Use of Radar Observations in Hydrological and NWP Models. Currently he is chair of the SRNWP (Short-Range NWP Programme) Expert Team on Data Assimilation and Use of Observations, a group organised by collaborating European Meteorological Services.



### **Participant no. 3: University Rovira i Virgili (URV), Spain**

The new Centre on Climate Change (C3), set recently up by University Rovira i Virgili (URV) in support of the work carried out by the former URV's Climate Change Research Group (CCRG), is chaired by **Dr. Manola Brunet** and supported by **Drs. Enric Aguilar and Javier Sigró**. It has a research experience and scientific background in the fields of climate data archaeology, instrumental climate reconstruction and analysis of climate variability and change and its forcing factors. C3's experience, through the former CCRG, is based on either the development of high-quality and homogenised climate datasets (at different temporal and spatial scales: i.e. at the daily scale in Brunet *et al.*, 2006) and on the assessments of climate variability and change (both on the mean and extreme states of the climate: i.e. Brunet *et al.*, 2007a,b). Its expertise in the field of the development of high-quality climate datasets is also endorsed by the contribution of its members to both normative World Meteorological/World Climate Data Monitoring Programme (WMO/WCDMP) guides on quality control and homogeneity (Aguilar *et al.*, 2003) and on the development of daily adjusted temperature records (Brunet *et al.*, 2008). The C3's has also contributed to the development and analysis of high-quality controlled and homogeneous climate datasets for the Globe, Europe, America, Africa, Middle East and Spain, with which has contributed to document changes in the mean and extreme states of the climate at regional and global scales for the last two IPCC-WGI reports (i.e. Alexander *et al.*, 2006; Ansell *et al.*, 2006; Moberg *et al.*, 2006). URV has been leading and taking part in several European and national research projects, like EMULATE at the European scale. URV has strong links and active involvement with international bodies and their activities, such as WMO/CCL, UNFCCC and GCOS.

Manola Brunet is currently co-chairing WMO/CCL OPAG II: Monitoring and Analysis of Climatic Variability and Change, she is member of the CCL Implementation and Coordination Team and is also co-chairing the WMO-MEDARE Initiative ([www.omm.urv.cat/MEDARE](http://www.omm.urv.cat/MEDARE)), which bring together climatologists from most of the GMR-NMHSs with scientists from universities, research centres and international bodies and projects with the main objective is being to develop, consolidate and progress climate data and metadata rescue activities across the GMR. She is also an UNFCCC/SBSTA international expert on data and observations in support of the Nairobi work programme on impacts, vulnerability and adaptation to climate change. Finally, Manola Brunet had an active role in the development of the GCOS Regional Action Plan for the Mediterranean Basin.

Two members of the C3, Enric Aguilar and Manola Brunet, have been actively involved in regional training activities addressed to the staff of NMHSs in RA I, II, III and IV organized by the Expert Team on Climate Change Detection and Indices (ETCCDI) for WMO Regions I, II, III and IV focused on analysis of climatic extremes (i.e. Aguilar *et al.*, 2005; Aguilar *et al.*, 2008; Peterson *et al.*, 2007; Zhang *et al.*, 2005).

#### **Participant no. 4: National Meteorological Administration (NMA-RO), Romania**

The National Meteorological Administration (NMA-RO) represents the national authority in the field of meteorology in Romania acting within the Ministry of Environment and Sustainable Development. NMA-RO is the owner and unique administrator of the meteorological, climatological and aerological Romanian databases. NMA-RO coordinates the National Meteorological Observation Network, which consist of: 162 meteorological stations, 2 aerological stations, 8 radar centers, 60 agrometeorological stations and 8 actinometrical stations. Our research activity is focused on climate variability and change at the regional scale and climate predictability. The main research topics are: analysis of the main characteristics of climate variability over Romania using long-term observations (trends, shifts, extreme events), connection between Romanian climate and large-scale phenomena (e.g. the North Atlantic Oscillation, Atlantic Multidecadal Oscillation), projection of global climate change on local scale using statistical and dynamical downscaling models, validation of global/regional climate models on large-scale and regional scale.

**Dr. Roxana Bojariu** is leading the Climate Research Group at the NMA-RO. Her expertise is in the field of climate variability and change and associated impacts. She has been involved in European projects such as EuroGLOBEC (European global ocean ecosystem dynamics), FP6 DYNAMITE (Understanding the Dynamics of the Coupled System), FP6 IPY-CARE (Climate of the Arctic and its role for Europe – an European component of the International Polar Year), FP6 CECILIA (Central and Eastern Europe Climate Change Impact and Vulnerability Assessment), FP7 METAFOR (Common Metadata for Climate Modelling Digital Repositories) and in international ones (e.g. Small Pelagic Fishes and Climate Change). She was lead author of the chapter on observations in the IPCC-WGI-AR4 and Review Editor of the AR4 Synthesis Report.

**Dr. Mihaela Caian**, senior research scientist, is leading the Numerical Modeling group and has more than 15 yr experience in numerical modeling for meteorology. She started the regional climate modeling at NMA-RO and participated in the construction and development of ALADIN and RegCM model for Romania. She has been leading or was involved in several international projects for the NMA-RO such as LIFE ASSURE, LIFE AIRFORALL.

**Dr. Sorin Cheval** is the Deputy Head of the Department of Climatology of NMA-RO. His expertise focuses on urban climate, climatic hazards, and GIS and remote sensing applications in climatology. He is a member in the Management Committee of the COST Action HOME. Sorin Cheval was a Fulbright grantee at the Hazards Research Lab, University of South Carolina, USA (2001-2002), and he has been involved in several national and international research projects. He is conducting the research project “The Climatic Atlas of Romania” sponsored by the Romanian Ministry of Environment (2007-2009).

**Dr. Constanta Boroneant** is senior research scientist in the Climate Research Group. Her expertise encompasses homogenization of climatological data, regional aspects of natural climate variability, extreme events, anthropogenic influence on climate, GCM and RegCM validation. She has been involved in the STARDEX project during a postdoc position (2004) at Institut Non Linéaire de Nice-CNRS and in several national climate research projects on climate change and its impacts. She has been acted as expert reviewer for WGII contribution to IPCC-TAR and to WGI and WGII or IPCC-AR4. Presently she is leading the NMA-RO research team in the FP6 project CECILIA. She is a member in the

Management Committee of the COST-Action HOME. She is member of Expert Team on Observing Requirements and Standards for Climate – OPAG I – Climate Data and Data Management, Commission of Climatology of WMO.

**Dr. Aristita Busuioc** is leading the Climatology Department and has a long experience in statistical analysis, her scientific interest being related to the statistical downscaling models. As expert in statistical downscaling models, she was involved in international projects such as the US Country Study Program, the Swedish regional climate modelling program, STARDEX and ENSEMBLES. She was lead author of chapter on regional projection in the IPCC-WGI-AR4 and the IPCC governmental focal point for Romania.

## **Participant no. 5: Meteo Swiss (MS), Switzerland**

The Federal Institute of Meteorology and Climatology MeteoSwiss (MS), is the governmental centre for information on weather and climate in Switzerland. It operates the national observation networks, issues weather forecasts and warnings, provides generic and tailor-made datasets for customers, and conducts research on themes from now-casting to climate prediction. Weather and climate in the Alpine region is one of its core competences. MeteoSwiss hosts the national GCOS office, it is currently presiding WMO RA VI Europe, and is national contact point to inter-national institutions (WMO, ECMWF, EUMETNET etc.). MeteoSwiss participates in the Swiss National Centre for Competence in Research on Climate (NCCR-Climat), in the EU FP6 project ENSEMBLES, in the CM-SAF and in several COST actions. In its research MeteoSwiss collaborates with academia (e.g. ETH Zurich), with other governmental offices (e.g. hydrology) and the private sector (e.g. reinsurance).

**PD Dr. Christof Appenzeller**, is head of the section Climate Services at MeteoSwiss (35 collaborators) and senior lecturer at ETH Zürich. He has long-standing research experience in the analysis and prediction of the atmosphere-ocean-cryosphere system and was PI of several research projects on climate variability and climate risk management. He is author of numerous papers, including Science and Nature, and is governmental representative in various commissions (ECSN, WMO and IPCC).

**Dr. Mischa Croci-Maspoli**, is head of the Climate Information Group. His research experience covers the dynamics of large-scale atmospheric flow and inter- and intraseasonal climate variability. He was a Co-PI of the ETH contribution to the Swiss NCCR-Climat project and is currently a member of the management committee of the COST ES0601 Action. In EURO4M he will collaborate with **Dr. Simon Scherrer**. Simon Scherrer's expertise encompasses climate variability and change, atmospheric predictability, dynamics and physics of the atmosphere and the hydrological cycle and data quality management in climate monitoring. He was a contributing author to the IPCC-AR4.

**Dr. Mark A. Liniger**, is head of the Climate Analysis Group, and as such responsible for several ongoing research projects on past, present and future climate. He has research experience in upper tropospheric and stratospheric transport, mid-latitude dynamics and intra-seasonal variability. His latest work focuses on the verification and application of dynamical seasonal forecasts, climate risk management and statistical data analysis. He is Co-PI of the MeteoSwiss contribution to the Swiss NCCR-Climat.

**PD Dr. Christoph Frei**, is senior scientist in the Climate Analysis Group. He has long-standing research experience in spatial climate analysis, evaluation of climate models, application of extreme value statistics in climatology, and climate change downscaling. He compiled a pan-Alpine rain-gauge dataset and derived widely used precipitation analyses. He was Co-PI in EU research projects (e.g. PRUDENCE, STARDEX), was a contributing author to IPCC-AR4, and is a lecturer at ETH Zürich. In EURO4M he will collaborate closely with **Dr. Reto Stöckli**, who is responsible for MeteoSwiss activities in CM-SAF. Reto Stöckli has many years of research experience in the analysis of satellite datasets, in land surface modelling and in radiation. He was also active in visualizing satellite data for public outreach and earth science education.

#### **Participant no. 6: Deutscher Wetterdienst (DWD), Germany**

The CM-SAF, which is under the guidance of the German Weather Service (DWD), is dedicated to the high-quality long-term monitoring of the climate systems state and variability, partly on the regional level. It supports the analysis and diagnosis of climate parameters in order to detect and understand changes in the climate system. The SAF, furthermore, serves the modelling of the atmospheric system as well as planning and management purposes. Utilising specialist expertise from the Member States, SAFs are dedicated centres of excellence for processing satellite data and form an integral part of the distributed EUMETSAT Application Ground Segment. Each SAF is led by a consortium of organisations under the guidance of a National Meteorological Service. The research, data and services provided by the SAFs complement the standard meteorological products delivered by EUMETSAT's central facilities in Darmstadt, Germany. EUMETSAT supervises and coordinates the overall activities of the SAF network and the integration of the SAFs into the various operations within the EUMETSAT Application Ground Segment.

**Dr. Richard Müller** is head of the radiation group of CM-SAF. Richard Müller has excellent expertises in different fields of atmospheric research, such as meso-scale modelling of wind fields, modelling, analysis and interpretation of stratospheric processes, radiative transfer modelling and development of radiation retrieval schemes. The main focus of his work within CM-SAF is the generation and analysis of radiation and cloud climatologies. He has published several articles in peer-reviewed journals. The expertise of Richard Müller is supplemented by other CM-SAF scientists, experts in climate monitoring and retrieval of atmospheric parameters. They have liaisons with GCOS and GEWEX, e.g. the scientific head of the CM-SAF. **Dr. Jörg Schulz** is member of the GEWEX radiation panel. In addition, Richard Müller studied pedagogics, with special focus on the adaptation of Paulo Freires concepts to “modern education” in natural sciences. Richard Müller has several years of training and teaching experience.

**Dr. Tobias Fuchs**, head of the WMO Global Precipitation Climatology Centre (GPCC) also participates in this project. The GPCC is operated by DWD. The Centre holds the world-wide largest global rain gauge database for monthly precipitation. Tobias Fuchs and the scientific staff members of GPCC are experienced in data processing, quality control, gridding and statistical analysis. The GPCC co-operates with NOAA and NASA within the GPCP on the merging of rain gauge and satellite-based observations. The Centre also co-operates with the Global Runoff Data Centre in water budget studies. Tobias Fuchs is Member of the GCOS Atmospheric Observation Panel for Climate (AOPC) and of the WMO Region VI Working Group for Hydrology.

## **Participant no. 7: Swedish Meteorological and Hydrological Institute (SMHI), Sweden**

The Swedish Meteorological and Hydrological Institute (SMHI), is a governmental institute under the auspices of the Swedish Ministry of the Environment. With expertise in meteorology, hydrology and oceanography, a well developed international network and access to advanced models, SMHI offers services and products to promote efficiency, safety and sustainable development in various areas of society. Vast quantities of data are gathered continuously in real time from land-based weather stations, balloons, ships, buoys, aircrafts, weather radars, satellites and lightning localisation systems. All information from SMHI's and other international observation systems is managed using powerful computers, and based on these observations highly advanced numerical models and statistical methods are used in real time to carry out analyses and forecasts which form the basis of further work. This includes air pollution forecasts and nuclear emergency preparedness on different scales. In addition, high-quality gridded analyses with high resolution in space and time are performed in non real time, covering Europe or parts of Europe and including meteorological, climate and environmental variables.

SMHI's products are presented and disseminated using state-of-the-art IT-technology, serving the differing needs and requirements of its large community of users from both the public and the commercial sector. SMHI's expertise is used in extensive analyses, climate studies and research. SMHI has approximately 550 employees. It has seven departments and its management system has been certified under the quality standards ISO 9001 and ISO 14001.

SMHI has a most keen interest on the European collaboration and participates enthusiastically in the ECMWF, HIRLAM and the EUMETNET activities. **Prof Nils Gustafsson** has been at the forefront of data assimilations since the early sixties. Within the HIRLAM consortium he has lead the development and implementation of not only the 3-dimensional variational analysis (3D-VAR) but also the full 4-dimensional version (4D-VAR). **Dr Magnus Lindskog** is the other SMHI expert on 3D-VAR and has been working closely with Nils Gustafsson and other colleagues in several of the HIRLAM countries. Magnus Lindskog has particular insight in the use and quality control of observations inside the variational assimilation. **Ulf Andrae** has a lot of modelling and actual reanalysis experience. During the first years of this century he developed and ran the BALTEX (Fortelius *et al.*, 2002) reanalysis for the areas around the Baltic catchment and made a lot of diagnostic computations of balances for the basin. **Per Undén** has extensive experience of data assimilation at ECMWF during earlier years and at SMHI he has 6 years of experience of leading and managing the HIRLAM-5 and HIRLAM-6 research programmes involving well over 50 scientists in 8 countries (Undén *et al.*, 2002). SMHI has played a strong role in the ECMWF reanalysis activities (**Dr Per Kållberg**) and presently leads the EUMETNET-ECSN Showcase EUROGRID (**Christer Persson**). SMHI continues investing heavily on the HIRLAM/ALADIN collaboration, not least on variational data assimilation. Furthermore, SMHI devotes considerable effort on the advanced mesoscale analysis system, MESAN, (**Dr Tomas Landelius and Anna Jansson**) both for Sweden and on the scale of Europe. **Assoc. Prof Lars Bärring**, formerly head of the Rossby Centre, the climate research and modelling unit of SMHI, has 20+ years experience in research on climate variability and experience from several European projects (e.g., ADVICE, WEELS, MICE, PRUDENCE, ESPON, ENSEMBLES).

Tomas Landelius, Nils Gustafsson, Per Kållberg, and Magnus Lindskog together with Per Undén, Ulf Andrae, Christer Persson and Anna Jansson will be key participants in WP2.

**Participant no. 8: Climatic Research Unit, University of East Anglia (UEA), United Kingdom**

The Climatic Research Unit (CRU) at the University of East Anglia (UEA) has over 35 years of extensive experience in the analysis of climate data. CRU jointly produces the well-known datasets of global surface temperatures (HadCRUT3 and CRUTEM3 - see [www.cru.uea.ac.uk/cru/data/temperature](http://www.cru.uea.ac.uk/cru/data/temperature)), as well as many other climatic variables ([www.cru.uea.ac.uk/data](http://www.cru.uea.ac.uk/data)). CRU has been involved in a number of studies on the analysis of long instrumental records with particular emphasis on extremes, as well as the development of long homogeneous series not only within Britain, but also in other parts of Europe. UEA were actively involved in the recent ALP-IMP and EMULATE projects and are involved in ENSEMBLES. The EMULATE daily pressure dataset (EMSLP) will be particularly useful for the analysis of changes in storminess and in storm tracks. CRU/UEA has also been involved in a recent assessment of radiosonde data and has experience of using the Integrated Surface Hourly (ISH) dataset. Both these datasets are likely to significantly improve the conventional input data used by the ERA-40 Reanalyses.

**Prof. Phil Jones** is Director of the Climatic Research Unit at UEA. He is also a member of the Atmospheric Observations Panel (AOPC) for GCOS and was one of the two coordinating lead authors on the Chapter on Atmospheric Observations of the 2007 IPCC report. Within the AOPC, he chairs the Advisory Group for the GCOS Upper Air and Surface Networks (the GUAN and GSN). This group assesses the performance of these two key networks and suggests improvements and changes to NMHSs as well as approving or not suggestions for network improvements made by NMHSs. He coordinated the EMULATE project and has been involved in numerous other EU projects including (IMPROVE, STARDEX, ALP-IMP) and is currently involved in ENSEMBLES, ECOCHANGE and CLARIS-LPB. He has over 30 yr of experience in the climate change field. He was awarded the Hans Oeschger medal from the European Geophysical Society in 2002 for work in paleoclimatology. Also in that year he was awarded the International Journal of Climatology prize of the Royal Meteorological Society for papers published in the past five years in the International Journal of Climatology. He is recognised as one of the top 0.5% of highly-cited researchers in the Geosciences field by the ISI (the institute in the US that maintains the Web of Science, where publications and citations are monitored). In 2007 he was awarded a fellowship by the American Meteorological Society. He has worked with most of the partners within the present proposal.

**David Lister** has been in CRU since the mid-1990s. He has worked on a number of EU-supported projects (including EMULATE and ENSEMBLES) as well as several for the Environment Agency in the UK as well as a number of consultancy type projects. He is adept at data handling, particularly when comparing observations with climate model output.

**Ian Harris** has been in CRU since the late-1990s. He has worked on a number of EU-supported projects (including HOLSMEER and one on the effects of climate change on historic building across Europe) as well as a number of NERC projects within the UK. He is currently working on updating the CRU High-resolution datasets. He is also adept at data handling, and will be well-suited to the software needed to read the NCEP/NCAR Reanalysis input as well as the ISH data.

### **Participant no. 9: Météo France (MF), France**

Météo France (MF) is the national research and information centre for weather and climate. The meteorological research is a major part of the activity of MF. With 250 researchers, MF plays a leading role in the international community, especially in the fields of climate research and atmospheric modelling. The NWP models, developed jointly with ECMWF for the global model ARPEGE-IFS and the ALADIN Group for a small scale numerical limited area model ALADIN and recently for the NH-model AROME. MF owns and maintains a climatological database which archives data acquired by more than 1300 automatic stations and over 3200 weather stations, some of which go back to 1850. This dense observation network allows statistical studies for research purposes and the elaboration of decision support tools for various sectors of the economy.

MF participates actively in several European collaborations, including HIRLAM, EUMETNET, GMES, and ENSEMBLES.

**Eric Bazile** is senior scientist at the research center of MF (CNRM-GAME) since 1991. He has several years of experience in the fields of surface data assimilation, surface and boundary layer parameterization for NWP. He has been involved and collaborates with both the HIRLAM and ALADIN consortia since 1993, in particular during the development and the operational implementation of the soil moisture assimilation and the ISBA scheme. He is member of the coordinating group in the NetFAM project (Nordic Network on Fine-scale Atmospheric Modelling).

**Dr Jean-Francois Mahfouf** is senior scientist. He worked at CNRM-GAME from 1988 to 1994 with main activities on land surface processes for NWP and climate modelling. From 1995 to 2002, he performed research at ECMWF on global variational data assimilation and on land surface analysis. He worked at Environment Canada from 2002 to 2006 on mesoscale data assimilation of soil and precipitations. In 2006, he moved back to CNRM with main activities on meso-scale data assimilation of soil and boundary layer observations. He is a Member of the Royal Meteorological Society since 2000.

**Eric Martin** is head of the CNRM-GAME team involved in atmosphere-surface-hydrology interactions research. This team has used and validated the SAFRAN analysis system at the scale of France. Eric Martin has a long experience in surface processes modelling, including snow cover modelling. He will be involved in the definition and the validation of the new analysis system by comparison to the present SAFRAN analysis. He will also assess the impact of the new analysis on the modelled surface fluxes.

**Dr Fabienne Rousset-Regimbeau** has worked at CNRM-GAME from 2003 to 2007, with activities on land surface processes, hydro-meteorological modelling and hydrological ensemble predictions (HEPEX experiment). Since 2007 she has been working in the team in charge of performing SAFRAN meteorological analyses, both in real-time operational and retrospective contexts. Since 2007, she has also been contributing to the H-SAF project (Satellite Application Facility on Support to Operational Hydrology and Water Management).

**Laurent Franchisteguy** has worked at CNRM from 1999 to 2004 with activities on land surface processes and particularly on retrieval of surface properties using remote sensing (contribution to both CYCLOPES FP5 and Land-SAF projects). From 2004, he has been working in the team in charge of performing SAFRAN meteorological analyses, both in real-time operational and retrospective contexts. Since 2005, he has also been contributing to the H-SAF project.



**Marie-Hélène Théron** is deputy-director of the Climate Data Management team at MF. She occupied various positions in the National Meteorological Service. She was namely involved in data monitoring and control. In her present affectation, Marie-Hélène Théron is in charge of the data collection, climatological control and validation. She is heavily involved in the French national coordination mechanisms in charge of data management.

**Pierre Lassègues** is senior engineer in the Climate Data Management team at MF. He occupied various positions in the National Meteorological Service. Since 1998, Pierre Lassègues is in charge of the data management and data bases, especially for model outputs. He is specialized in valorisation of different sources of climatological information, namely reanalysis.

## 2.3 Consortium as a whole

A strong and enthusiastic consortium has been built to develop the capacity for, and deliver the best possible and most complete (gridded) climate change time series and monitoring services for Europe. Core participants are listed in the table on page 2. These 9 partners provide the necessary expertise for regional observation datasets, regional reanalysis and user-oriented climate change monitoring products and services.

Collectively, this is a very strong group of experienced researchers with proven expertise in handling complex projects. Their complementarity (see also the participant descriptions) ensures that the whole spectrum of climate change monitoring is covered. The geographical spread of EURO4M participants is well distributed across the European domain, and this is enhanced further through linkages with other European organisations, institutions and agencies.

No.	Third party participant organization name	Contact
1	European Environment Agency (EEA)	Thomas Klein
2	European Centre for Medium-Range Weather Forecasts (ECMWF)	Philippe Bougeault
3	Network of European Meteorological Services (EUMETNET)	Steve Noyes
4	European Climate Support Network of EUMETNET (ECSN)	Aryan van Engelen
5	Global Climate Observing System (GCOS)	Stephan Bojinski
6	WMO Commission for Climatology (CCL)	Pierre Bessemoulin
7	National Oceanic and Atmospheric Administration - National Climatic Data Center (NCDC), US	Russell Vose
8	National Oceanic and Atmospheric Administration - Earth System Research Laboratory (ESRL), US	Gil Compo
9	Queensland Climate Change Centre of Excellence (QCCCE), Australia	Jozef Syktus
10	Environment Canada	Francis Zwiers

Table 2.3a. Details of European and international third party participants in EURO4M.

Table 2.3a lists the key stakeholders that have already indicated their commitment and interest in the project. These are linked to the EURO4M consortium as so-called third party participants. They are representatives of the wider network of interactions across the globe and will be involved in the General Assemblies and/or Advisory Board (see Section 2.1) and through the organization of strategic workshops. Together, they make the consortium even stronger.

**Key third party participants who have indicated that they strongly endorse the project (see the support letters in Appendix B) and with whom special arrangements will be set up (through letters of agreement or memorandums of understanding) include EEA, ECMWF, EUMETNET, and GCOS:**

- EEA is expected to play an important coordination role in the *in situ* component of GMES (EC, 2008), which EURO4M will take into account. EEA have a strong interest in the availability of multi-decadal and high-resolution climate monitoring information and data beyond current state-of-the-art. They will act as one of the principal customers for CIBs, which feed into their environmental assessment reports for Europe (see Figure 1.1 a,b).
- ECMWF is suggested within GMES as a technical centre that coordinates the Atmosphere network and ensures the provision of operational atmosphere services (EC, 2008). Therefore, a clear link between ECMWF and EURO4M will exist. ECMWF will also provide necessary boundaries and observations for the regional reanalysis in EURO4M, whereas EURO4M will deliver valuable input for future ECMWF global reanalysis.
- EUMETNET in their European Climate Support Network (ECSN) aims at providing sustainable high quality and climate reference services, tailored to the users needs. EURO4M will work together with EUMETNET-ECSN on the production of gridded climate data sets based on dense networks of station observations, particularly in the EUROGRID proposal which is in preparation by ECSN members for the EUMETNET Council (as a follow-up to S-EUROGRID).
- GCOS requirements for global climate monitoring are largely met with EURO4M in place. EURO4M is positioned to form the implementation component of the atmospheric sections of GCOS. The focus on extremes and weather related hazards in EURO4M even provides more detail than is strictly necessary for global climate monitoring.

The ambitious scope of the project means EURO4M must be implemented as a collaborative project. To span this entire range demands a multi-disciplinary critical mass with support from operational (national) meteorological agencies, professional data centres, International and European project teams, universities, and climate analysis centres. EURO4M needs a consortium linking EU Member, Accession and Candidate State institutions, and links to international initiatives, which have scientific expertise and ongoing responsibilities for atmospheric data networks, databases and operational monitoring. Only then will the project ensure that scientists, decision makers and application communities have the best data products, indicator abstractions and support for their particular needs.

A good illustration of the need for integrated efforts is the summer of 2005 with wet extremes (floods) in Romania and the Alpine areas and dry extremes (forest fires) in Portugal and Spain. High quality integrated monitoring products for these events were hard to obtain online, in particular products that use all available data sources to place these events in the context of long-term climate variability and change.

**The EURO4M consortium contains Europe's leading expertise on the monitoring of climate change.** Four participants have been heavily involved in the production of the IPCC-WGI-AR4 as lead author (3×) or coordinating lead author (1×). Therefore, the group has considerable experience in what it takes from a scientific point of view to gather the information needed to prepare high quality policy relevant assessments. Many

of the participants have already successfully co-operated in a number of EU projects, and have a well-developed capacity to collaborate effectively and reach important scientific goals together. However, this project forms a unique opportunity to integrate the entire community for the first time in order to achieve major advances in the key science issues in harmonizing the monitoring of climate change and extremes.

In summary, the consortium possesses the scientific and technical experience to administer and implement a project of the size and complexity of the proposed project. A number of EURO4M core participants have experience in the coordination of EU projects, in which WP-leaders have played major roles (leading WPs). This will directly benefit the current consortium. EURO4M consortium participants are also strategically placed to liaise with a wide range of international and national bodies, which will strongly leverage the EURO4M resources. **In particular, all European NMHSs are linked to EURO4M via its close association with ECMWF, EUMETNET, EUMETSAT, and their joint European Meteorological Infrastructure (EMI).** These linkages, and the overall expertise and experience of the participants, will ensure the success of EURO4M, as well as providing a mechanism for feeding the project results into the international domain.

***Sub-contracting***

None of the EURO4M participants makes use of sub-contracting.

***Potential new participants***

At the moment, it is not foreseen that any new participants will join EURO4M. However, it may become necessary during the course of the project that a third party participant organization joins the EURO4M core consortium.

**2.4 Resources to be committed**

The substantial resources requested overall for EURO4M reflect the ambitious scope of the project. Even with a budget of this magnitude, partner institutions will need to contribute significant additional resources. More importantly, **EURO4M needs to use to the largest extent possible existing capacities.** The expertise and experience of the participants in each WP, together with the degree of European and international co-operation, make the work feasible within the level of resources available. **The total grant requested from the EU is 4 million Euros (M€),** broken down as shown in Table 2.4a.

WP	Name	K€
1	Regional observation datasets	1470
2	Regional reanalysis	1710
3	User-oriented information and climate change products	540
4	Project management, coordination and sustainability	280

Table 2.4a. Breakdown of total grant requested amongst EURO4M WPs.

Most of the budget is used to hire personnel; a small fraction is for computing and travel. From the total budget of 4 M€ roughly 80% will be spent on science to make available high quality ECV datasets in WP1 and WP2. The other 20% will be spent on integrated products (at the required level of aggregation and processing) and management. We believe this breakdown is justified, because European climate change monitoring requires a firm scientific basis. Focusing on existing datasets and systems only, would result in products and services that are far from the forefront of scientific developments. About 7% of the overall budget is dedicated to management. This small amount implies that a large part of the support needed will come from either the Coordinators home institute (e.g. for hosting the EURO4M website) and from the home institutes of the WP-leaders.

### **Data storage and dissemination**

A huge amount of observational and reanalysis data will be produced in WP1 and WP2. These data will be analysed within the project and used for the multi-purpose products developed in WP3. WP3 will make an inventory of raw data needs by users outside the project as part of the CLT activities. We anticipate the need for EURO4M data after the end of the project. Therefore, the main datasets will be secured for at least 5 yr after the project ends. This is possible with the allocated resources only by virtue of exploiting existing infrastructure to the full, and capitalising on the benefits associated with an adherence to standards and interoperability.

Data sharing within EURO4M will follow a service-oriented approach adopting common standards on metadata, data models, and network services as described by INSPIRE and the GEOSS Architecture and Data Committee. **MO** leads the special group within EUMETNET called INSPIMET, which looks after the implementation rules of INSPIRE for meteorological and climatological data and services. Following these rules will ensure global connectivity and interoperability, also providing appropriate links to the WMO Information System (WIS). **The INSPIRE-compliant infrastructure will enable considerable data-sharing efficiencies to be realised between EURO4M and the various GMES services requiring access to climate-related data.** WP3 will describe the infrastructure that is used for the datasets produced in EURO4M.

### **Justification of resources by WP**

#### **WP1: Regional observation datasets**

The range of activities to be undertaken in WP1 has close linkages and co-operation with ongoing European and international initiatives and infrastructures. We rely on a large number of facilities, instruments and services owned and operated at national, regional and international levels inside and outside the EU. The work in this WP is designed in a way that recognizes that the *in situ* infrastructure is developed and maintained by Member States and should remain their responsibility (EC, 2008). The funding is used for pan-European integration, which contributes to globally co-ordinated data collection and exchange. The range of expertise in WP1 embraces both long experience in climate monitoring and dataset development of specific ECVs and ongoing involvements on European and international scientific bodies that have developed the current best practises and protocols for such activities. This allows for maximum optimal utilisation of the resources available. The data products will be stored and disseminated using existing data infrastructures, including the KNMI Climate Explorer ([climexp.knmi.nl](http://climexp.knmi.nl)) and CM-SAF web user service.

## **WP2: Regional reanalysis**

This WP gains cost effectiveness by building on developments already funded (“leverage”). In particular the work packages WP2.1-3 build on the NWP data assimilation system which are operational at MO, SMHI and MF. Similar use of well-tried operational systems has been a characteristic of most successful reanalysis projects. The infrastructure that needs to be exploited to the full in WP2 is partly from ECMWF. ECMWF is the leading NWP centre in the world, resulting from a 30 yr investment of its 25 European Member and Cooperating States. Its capability to undertake global reanalysis derives both from this core investment and from the external investments that have been made in past reanalysis projects by the European Union and supporting institutions outside Europe.

WP2 will use boundary conditions and the observational database infrastructure and expertise on observational bias and quality from the global ERA reanalysis. In particular, the intermediate reanalyses in WP2.2 can be undertaken by the project only because they exploit the ERA-40 and the new ERA-Interim reanalysis being undertaken by ECMWF and the extensive infrastructure and experience built up as a result of ECMWF’s operational and earlier reanalysis activities. Reanalysis is the tool that is generally seen as the ultimate tool for integration of all climate change ECV data. As indicated in Section 1.1, it will take some time before this ideal situation is reached, but the EURO4M investment will strongly contribute to making this happen.

## **WP3: User-oriented information and climate change products**

This WP will introduce, develop and implement new CIBs that provide integrated climate change information to different user groups. The CIBs form the main outcome of EURO4M and may develop into a future operational GMES service on climate change monitoring that directly addresses the user needs. The share of the budget dedicated to this WP is therefore fully justified. It can even be considered low, but WP3 will fully profit and build upon the work in the other WPs and the extensive national and international competence amongst the EURO4M participants. The user interaction aspects in this WP will be supplemented by work already performed within the CM-SAF and the user liaison work already in place at a national level. All of the participants in WP3 have means of asking user feedback in place at a national level. These networks will form the basis for the European CLT, together with other international networks, such as the ERA-Net CIRCLE. The fact that this WP receives the lowest share of the budget reflects that the CIB and CLT work is less labour intensive compared with the scientific work in WP1 and WP2.

## **WP4: Project management, coordination and sustainability**

The provision of a separate WP for project management, coordination and sustainability reflects the recognition that a project of this scale and ambition must be very efficiently managed if the project is to fully deliver its extensive plans. KNMI will coordinate the EURO4M project. This will involve organising a kick-off meeting for the project, organising EURO4M general assembly and management board meetings, communicating regularly with the EC, monitoring the progress of the project, and submitting management progress reports and scientific and technical progress reports to the EC. The small management budget for WP-leaders is needed to guarantee that the work flow and reporting from each individual WP evolves according to plan.

## 3. Impact

### 3.1 Expected impacts listed in the work programme

#### *Expected strategic impact of the proposed project*

EURO4M will contribute to establishing a data archive of systematic observational data related to the climate system. The climate change time series will be based on the optimal combination of regional observation datasets of Essential Climate Variables (ECVs) and model based regional reanalysis. A continuous record of ECVs will be developed, coherent with UNFCCC requirements. EURO4M will contribute to the consistency of such a dataset, as well as to a sustainable and transparent access to such data for global climate scientific and operational communities. This paves the way for a sustainable provision compliant with the requirements of climate analysis communities.

The high costs and joint responsibilities of climate monitoring have always favoured international co-operation, which helps to avoid duplication and promotes sharing of information. The World Weather Watch (WWW) of the WMO is a very positive example of successful international co-operation. Unfortunately, many national climate monitoring systems still operate independently, exchanging little or no data or information. Significant temporal and spatial gaps exist and therefore European integration is necessary. By providing a complete and accurate picture of the history of our atmosphere throughout the period of quantitative human observation, the regional reanalysis activity eventually provides basic input material for a large variety of activities in the domains of science, policy and applications.

In addition, many potential users see little of the data produced and are not offered data, information, products and services tailored to their needs. For instance, decision makers and policy-makers need information summaries in the form of indicators and indices, which are presently neither readily available nor based on sound scientific understanding and indisputable evidence. As illustrated in Figure 3.1a, aggregated solar energy maps are needed, rather than the underlying raw satellite data. Without these summaries it will be difficult to move into the sustainable path, where (according to the EU Sustainable Development Strategy) environmental protection goes hand in hand with economic prosperity and social cohesion.

Through the development of internationally recognised datasets, feedback and user interaction protocols, EURO4M will work to develop new high quality data products and services for the evaluation of severe climatic events that will aid both intermediate- and end-users, whether institutional, civil society organisations (e.g. the Red Cross/Red Crescent) or from the private sector (e.g. reinsurance). EURO4M will initiate and facilitate dialogues between scientists and end-users leading to a better understanding among scientists on the information needs of end-users and a better understanding of the end-users on how the available data-products can be understood. **The EURO4M system has the potential to evolve into a future GMES service on climate change monitoring that is fully complimentary and supporting the existing operational services.**

The results and outputs of EURO4M will also provide a new way to support climate-policy-related research at the national and local scale. National decision makers and local authorities will be able to utilise the state-of-the-art EURO4M data products and

services for their country or region as input to climate change assessments, and the formulation of adaptation and mitigation strategies. It is the longer multi-decadal time scale addressed in EURO4M that is needed for governments to minimize and adapt to the societal and environmental impacts of climate variability and change. European countries can directly use the results of the proposed project for their “national communications on climate change policies” which are a written requirement for the Conference of the Parties of the UNFCCC and include national GCOS implementation activities.

### Annual Solar Energy Europe




Figure 3.1a. Map of calculated solar energy in a typical year.  
Source: CM-SAF lead by DWD.

### ***Contribution to the scientific, technical, wider societal and policy objectives***

Long-term and reliable climate data are vital for detecting, understanding, predicting and responding to climate change and variability. The development of increasingly sophisticated climate models has reinforced the need for basic observations. However, as the demand for objective information grows, GEO and GCOS have indicated that our capacity to monitor climate and environmental changes and fluctuations is still largely inadequate.

Monitoring climate change and extremes at the multi-decadal and century time scale is currently not among the core GMES services. Ongoing activities in the GEO work plan also do not cover long-term climate observations and monitoring, even though GEO is a strong advocate for sustained and coordinated climate observing systems. EURO4M is positioned to form the implementation component of the atmospheric sections of GCOS and GEOSS. By taking on the climate section of the plan EURO4M will coordinate a monitoring framework essential for the success of GEOSS.



**EURO4M is crucial for establishing the climate change component of GMES and thus for the European contribution to the development and integration of the GEOSS.** A key system for climate change observations and information will be provided for the benefit of each of the nine societal benefit areas of GEO. The system integrates European climate change monitoring activities into the global picture. In this way, the project will contribute to forward European and international efforts in strategic policy areas, such as climate change and energy. Involvement of international partnerships will ensure proper know-how is taken into account, as well as avoiding duplication of efforts. This will enable Europe to play a pivotal role in the systematic monitoring of climate mandated by the UNFCCC.

EURO4M will form a core component of the European implementation of the atmospheric climate component of GCOS in support of the UNFCCC. The Implementation Plan for GCOS addresses the requirements identified in the Second Report on the Adequacy of the GCOS (GCOS, 2003), in particular the ECVs and associated climate products defined in the report. The UNFCCC Conference of the Parties endorsed the Implementation Plan at its 10th session in December 2004 through a formal decision which encourages Parties to strengthen their efforts to address the priorities identified.

Finally, within the context of the Kyoto Protocol, EURO4M will directly focus on the following Articles: 9.1) Provide the best available scientific information and assessment on climate change and its impacts; 10.c) Promote education and training, and increase public awareness of climate change; 10.d) Participate and co-operate in international scientific research programmes to reduce uncertainties related to the climate system and the adverse impacts of climate change.

### ***Contribution to standards***

The formation of the EURO4M consortium responds directly to the Commission's recommendations to mobilise all European resources through coordinating and networking efforts in Research and Technical Development areas related to climate change and environmental monitoring.

EURO4M will link EU and international system outputs, capitalising on the benefits associated with an adherence to standards (e.g. Open GIS Consortium developments, ISO 19100 series, Geographical Markup Language), interoperability (e.g. within INSPIRE and the WMO Information System (WIS)) and consolidation (see Section 2.4).

### ***Added-value in carrying out the work at a European level***

EURO4M will only work when carried out at the European level, building on existing national monitoring capacity and atmospheric observation programs to form a truly European integrated system capable of providing CIBs on extreme events that threaten Europe, and its citizens and interests worldwide. The development of such a system requires an unprecedented level of collaboration and coordination of multidisciplinary research, enhancement of operational activities within the climate research and monitoring community.

Although elements of such a system presently exist, the state of their development in the different European countries is uneven. More critically, access to data, information, and products by users is in many cases inadequate and difficult.

**A European-wide coordination will avoid unnecessary duplication and hence enable resources to be focussed more effectively. This will also result in more uniform, high quality and standardised products, which will be available to all European countries, reducing the geographical disparity between countries and regions for information and services available to intermediate- and end-users.** It will encourage the development of scientific research and analysis at the national level as well as at the EU level, allowing countries to make the most of the collaboration both in terms of data resources and R&D expertise. As a side-effect, improved access to products and services will encourage smaller nations to make a greater contribution to existing observation networks.

The enlargement of the European Union in 2004 gives the opportunity to enlarge the scientific co-operation. In fact, co-operation with East and Southeast European countries is essential to tackle topics in environmental research with a wider regional dimension such as climatic change and variability. The European scale ensures that sufficient resources are mobilised to address the goal and objectives of the work program and maintain pre-eminence within the international community.

### ***Account taken of other national or international research activities***

EURO4M is designed to build on existing capacity. As described in Section 1.1, the project fully benefits from ongoing national and international work on observational datasets and data assimilation. **EURO4M will link very strongly to the wider international community.** GMES cannot be successfully implemented without exchanging equivalent observational data through cooperation schemes, thereby sharing the cost of observation infrastructure with non-EU partners (EC, 2008).

EURO4M participants include members of the GEO Committees, GCOS Steering Committee, its Atmospheric Observation Panel for Climate (AOPC) and Oceanic Observation Panel for Climate (OOPC) specialist groups, WMO Open Programme Area Groups (OPAGs) and Expert Teams (some of them also lead by EURO4M participants), International research programme on Climate Variability and Predictability (CLIVAR) Expert Teams, the WCRP Observations and Assimilation Panel and the WCRP WG on Observational Data Sets for Reanalyses. EURO4M participants are also intimately involved in IPCC activities.

EURO4M will work with its international partners to collate and report on ECVs, including integrating existing data products and providing uncertainty estimates for each ECV. EURO4M will link with WMO programmes and improve the reporting of observations to international data centres. **EURO4M fits perfectly in the WCRP strategy for Coordinated Observation and Prediction of the Earth System (COPES; WMO, 2005).**

### ***Links to ongoing FP projects***

EURO4M links to ongoing FP6 and FP7 projects, including those developing prototype and pre-operational GMES services (such as GEOLAND2, MyOcean, MACC), and other related FP projects (such as ENSEMBLES, EURO-LIMPACS, CECILIA, CIRCE, etc.):

- EURO4M will complement GEOLAND2, which joins a range of previous GMES projects to deliver a range of geo-information services for the terrestrial sphere, and MyOcean, which will define and set up a concerted and integrated pan-European capacity for ocean monitoring and forecasting. EURO4M partners involved in GEOLAND2 and MyOcean will link the project results to these services.

They both require the long-term climate change monitoring information that EURO4M will deliver (in particular on extremes) as an input.

- Strong ties also exist with the MACC consortium lead by ECMWF. MACC (Monitoring Atmospheric Composition and Climate) combines the earlier projects PROMOTE and GEMS, which provide prototype atmospheric services for GMES. PROMOTE (PROtocol MOniToring for the GMES Service Element: Atmosphere) focuses on stratospheric ozone depletion, surface UV exposure, air quality, whereas GEMS (Global and regional Earth-system Atmosphere Monitoring using Satellite and *in situ* data) focuses on global distributions of atmospheric constituents important for climate. **Although responsible for the pre-operational atmospheric services from 2009 onwards, MACC does not include the longer (multi-decadal) time scales and many of the ECVs required for climate change monitoring that EURO4M will address.**
- EURO4M will build on the high-resolution gridded observational datasets for Europe developed in ENSEMBLES and work towards getting these datasets into an operational status. Our user-oriented work will benefit from collaboration with the EURO-LIMPACS (Integrated project to evaluate impacts of global change on European freshwater ecosystems) team, designed to assess the effects of future global change on Europe's freshwater ecosystems, as well as from CIRCE and CECILIA for collection and homogenisation of long daily temperature and precipitation data for all the Mediterranean countries and Eastern Europe.
- The work on extremes in WP3 will be performed through strong connections with the EU projects ENSEMBLES, CIRCE and CLARIS-LPB (A Europe-South America Network for Climate Change Assessment and Impact Studies). The work on extremes will also build on the results of earlier EU FP5 projects, like STARDEX, MICE (Modelling the Impact of Climate Extremes), EMULATE and ALP-IMP (Multi-centennial climate variability in the Alps based on Instrumental data, Model simulations and Proxy data), plus recent global initiatives from IPCC which many EURO4M participants have been actively involved with.

A clear link also exists with the global reanalyses FP7 proposal ERA-CLIM, which is in preparation by a consortium led by ECMWF. EURO4M is designed to be complementary to global reanalysis efforts, by concentrating on aspects where it can add value, such as a higher horizontal resolution (the ERA-CLIM proposal is suggesting 40km resolution), the assimilation of surface precipitation data, and delivering integrated products at the required level of aggregation and processing to respond to a wide range of users and downstream services.

### ***Links to other activities***

In an effort to further harmonize European climate change information, EURO4M will connect to the European Co-operation in the field of Scientific and Technical Research (COST-) Action ES0601 (HOME) on comparisons of methods for homogenisation of long instrumental climate records. New quality control and homogeneity testing of early instrumental observations, particularly from EU Accession and Candidate States and neighbouring countries will build on the data products generated by EU-funded and other projects, such as ECA&D, NORDKLIM (Nordic co-operation within climate activities), EuroClim (European Climate Change Monitoring and Prediction System), MedCLIVAR

(Mediterranean CLIVAR) and CLIWOC (Climate of the world's oceans), integrating these data with the holdings of a suite of international databases.

EURO<sub>4</sub>M will work with International Comprehensive Ocean-Atmosphere Dataset (ICOADS) initiatives to aid the digitisation of many millions of historical marine data in European archives being funded separately under projects such as ACRE. **EURO<sub>4</sub>M will link via ACRE to the 20th Century Reanalysis Project and other longer historical reanalyses.** With these reanalysis products, there are also the full linkages and “end-to-end” infrastructure which ACRE integrates in working with climate applications users.

EURO<sub>4</sub>M has already linked with the EUMETSAT-SAFs, in particular to assess user community needs and to contribute to the development of integrated products. The coordinator (at DWD) of the CM-SAF is participating in EURO<sub>4</sub>M.

EURO<sub>4</sub>M also links to MEDARE activities on data rescue and surface climate reconstruction for the GMR (under the umbrella of WMO; see Brunet and Kuglitsch, 2008), EUMETNET-ECSN for the remainder of Europe and EuroCryoClim for the Arctic. We will liaise with other recent initiatives for particular sub-regions such as the Alps and the Baltic Sea region. Examples are the project MOnitoring Climate variability and Change for an improved environmental and risk management in the Alpine space (MOCCA) and the Baltic Sea Experiment (BALTEX). The latter will seek to exploit and analyze the results of EURO<sub>4</sub>M to quantify the water and energy budgets over the Baltic Sea basin and evaluate and improve regional climate models. The good results in the spatialization tools for complex terrain achieved in the EUMETNET-ECSN HRT-GAR project constructing a High resolution Temperature Climatology for the Greater Alpine Region will also be taken into account. These kinds of products are useful to fill the existing gap between the present day observation datasets for Europe and what end users really need in terms of spatial resolution to support impact evaluation.

## 3.2 Dissemination and/or exploitation of project results, and management of intellectual property

### *Dissemination and/or exploitation plans*

EURO<sub>4</sub>M provides an exciting and unique opportunity to bring together existing meteorological services, universities, research institutions and programs in order to provide a seamless integration of climate change data and information flows and feedbacks. **The proposed Climate Information Bulletins (CIBs; see WP 3.1) and Climate Liaison Team (CLT; see WP 3.2) will provide a framework for a new generation of flexible products that foster better dialogue between data providers and data users, and integration of end-user requirements for environmental and societal applications.**

The effective dissemination strategy includes the identification of the target audience at regional and global levels by the CLT. An analysis of relevant interested stakeholders will be carried out and a list of all other relevant stakeholders will be produced. Then the most suitable communications programs and products (meetings, conferences, seminars, workshops, publications, brochures and leaflets) will be identified and a timeframe will be defined for their delivery, as specified in WP 3. In particular, the CIBs will be used to demonstrate the project outcomes to all participating and interested stakeholders. All project activities, related documents and results, published materials (including training

materials and reports) and events announcements will be regularly posted on the project website.

The knowledge and expertise developed in the project will be passed on to the young scientists and engineers in training through university professors involved who teach at their institutions and supervise PhD and graduate students. Chances will be provided for young scientists to be integrated into, and gain experience in, high level European research and international collaboration. National workshops will facilitate dialogue between the scientific institutes involved and representatives of operational sectors.

Specific communication activities, targeted at the general public, schools and the press, will be developed in order to stimulate interest in youth audiences, to attract future scientists in the broad fields of geo-sciences, and to increase public awareness about climate change and the risks of extreme weather events, leading to a better integration of these risks in the policies and programs of operational sectors. In addition, communication activities will also address educational outreach to the public with respect to basic knowledge about climate change.

### ***Educational links and capacity building***

EURO4M will foster better integration of research laboratories and operational meteorological agencies in education programs. The project will catalyse a program of visiting scientists and postdoctoral positions through existing institutions. Emphasis will focus on fostering interactions between existing, new, Accession and Candidate EU States and neighbouring countries.

In order to promote common approaches and develop consensus best practice for climate data, a series of workshops will be organised, and specialised collaborations will be sponsored on specific topics of relevance to the project. The CM-SAF at DWD has already established a training group, which is operating in an international network in cooperation with EUMETSAT. The involvement of universities in the consortium means that the EURO4M approach and data products will be incorporated in existing earth system education programs. EURO4M funding will support Masters, PhD and post-doctoral fellowships at some of the participant institutions.

**The consortium will also engender access to developing countries, which will be amongst the largest potential beneficiaries of international co-operation in earth observation.** The latter will be done through liaising with the series of worldwide workshops (Peterson and Manton, 2008; see also Klein Tank *et al.*, 2006) organised by the joint Expert Team on Climate Change Detection and Indices (ETCCDI) of the CCL, CLIVAR and JCOMM (Peterson *et al.*, 2001). The EURO4M Coordinator is currently also co-chair of this Expert Team. This will complement the GMES activities in the field of capacity building as discussed at the Lisbon meeting in December 2007 and contributes to the Nairobi work programme on impacts, vulnerability and adaptation to climate change of the UNFCCC. In addition, outreach activities on data rescue, preservation, digitization and homogenization techniques and procedures programmed jointly with the WMO MEDARE Initiative will also contribute to capacity building on these subjects in developing countries and Least Developed Countries.

## 4. Ethical issues

The participants in this project declare that there are no ethical issues relevant to this proposal. Nevertheless, the Coordinator and MB will continue to keep this aspect under review, and should any such issues arise in the future, they will be addressed and taken into account in due time.

## 5. Consideration of gender aspects

### *Gender Action plan*

Many of the participants in EURO<sub>4</sub>M have gender action plans at the institutional level as part of their commitment to gender equality. These include programmes to raise awareness of the issues involved in gender equality, commitments to family friendly work practices and career breaks, and provision of child-care facilities. Organisational initiatives to encourage gender equality enjoy high level backing within many partner institutes. For example, KNMI is a participant in the Dutch GAIA-network and as such actively involved in the EU EQUAL-II Priority for Participation project. MO have an ongoing equality training programme, coordinated at boardroom level, which is mandatory for all staff. For participants based at institutions with less detailed plans, the Gender Action plan in EURO<sub>4</sub>M will provide an important impulse.

One of the four WP-leaders of EURO<sub>4</sub>M is a woman, as are 2 of the lead scientists at the 9 participating institutes.

**Project Gender Committee:** The gender committee will actively promote the role of women at all levels within the project. It will be responsible for ensuring that the gender action plan is applied across the spectrum of WPs in the project, both in terms of internal communication of developments and progress via the project website, and communicating progress externally, via the annual gender action report. The committee will also be responsible for ensuring that the training and dissemination aspects of the project are female-friendly. The committee will consist of 3 members elected by all project participants on an annual basis, with the possibility of re-election.

**Annual Gender Action Report:** The report will document the extent to which actions promoting gender equality have been performed at the project level, and will chart the rates of female participation at all levels of the project.

**Recruitment of Female Researchers:** Recruitment of young, talented female researchers will be encouraged in EURO<sub>4</sub>M. Job advertisements will state the project's commitment to equality and to a family-friendly working environment and will explicitly encourage women to apply. The gender committee will liaise with national programmes in the production of suitable information material for schools.

**Consortium Agreement:** The Consortium Agreement governing the operation of the project will enforce the following minimum requirements on the participating institutions:

1. encouragement of applications by female researchers in job advertisement;
2. formal action to ensure that employees are properly informed about their parental rights and responsibilities;
3. encouragement of female coaching and mentoring schemes, and project management;
4. production of an annual report on the nature and utility/success of gender actions undertaken;

**Management Board:** The Management Board has been chosen to ensure that women are adequately represented at the highest organisational levels of the project and consists of seven people. From the four appointed, one is woman (25%). Whilst not approaching equality, this percentage is higher than that of women in senior positions in climate science generally, and gives women a significant say in how the project is organised and run.

## References

- Aguilar E., I. Auer, M. Brunet, T.C. Peterson, J. Wieringa, 2003: Guidelines on climate metadata and homogenization. WCDMP-No. 53, WMO-TD No. 1186. World Meteorological Organization, Geneva.
- Aguilar, E., T.C. Peterson, P. Ramirez Obando, R. Frutos, J.A. Retana, M. Solera, J. Soley, I. Gonzalez Garcia, R.M. Araujo, A. Rosa Santos, V.E. Valle, M. Brunet, L. Aguilar, L. Alvarez, M. Bautista, C. Castañon, L. Herrera, E. Ruano, J.J. Sinay, E. Sanchez, G.I. Hernandez Oviedo, F. Obed, J.E. Salgado, J.L. Vazquez, M. Baca, M. Gutierrez, C. Centella, J. Espinosa, D. Martinez, B. Olmedo, C.E. Ojeda Espinoza, R. Nuñez, M. Haylock, H. Benavides, and R. Mayorga, 2005: Changes in precipitation and temperature extremes in Central America and Northern South America. *J. Geophys. Res.*, **110**, D23107, DOI 10.1029/2005.
- Aguilar, E., A. Barry, M. Brunet, L. Ekang, A. Fernandes, M. Massoukina, J. Mbah, A. Mhanda, D.J. do Nascimento, T.C. Peterson, O. Thamba, M. Tomou, X. Zhang, 2008: Changes in temperature and precipitation extremes in Western Central Africa, Guinea Conakry and Zimbabwe, 1955-2006. *J. Geophys. Res.* (in press).
- Alexander, L.V., X. Zhang, T.C. Peterson, J. Caesar, G. Gleason, A. Klein Tank, M. Haylock, D. Collins, B. Trewin, F. Rahimzadeh, A. Tagipour, P. Ambenje, K. Rupa Kumar, J. Revadekar, G. Griffiths, L. Vincent, D. Stephenson, J. Burn, E. Aguilar, M. Brunet, M. Taylor, M. New, P. Zhai, M. Rusticucci, J.L. Vazquez-Aguirre, 2006: Global observed changes in daily climate extremes of temperature and precipitation. *J. Geophys. Res.*, **111**, D05109, doi:10.1029/2005JD006290.
- Alexandersson, H., *et al.*, 2000: Trends of storms in NW Europe derived from an updated pressure data set. *Clim. Res.*, **14**, 71-73.
- Allan, R.J. and T.J. Ansell, 2006: A new globally complete monthly historical gridded mean sea level pressure dataset (HadSLP2): 1850-2004. *J. Climate*, **19**, 5816-5842.
- Ansell, T., *et al.*, 2006: Daily mean sea level pressure reconstructions for the European - North Atlantic region for the period 1850-2003. *J. Climate*, **19**, 2717-2742.
- Auer I., *et al.*, 2005: A new instrumental precipitation dataset in the greater alpine region for the period 1800-2002. *Int. J. Climatol.*, **25**, 139-166.
- Babst, F., R.W. Mueller, R. Hollmann, 2008: Verification of NCEP reanalysis shortwave radiation with mesoscale remote sensing data. *Geosciences and Remote Sensing Letters*, **5**, 34-37.
- Bauer, P. and P. Schuessel, 1993: Rainfall, total water, ice water, and water vapour over sea from polarized microwave simulations and special sensor microwave/imager data., *J. Geophys. Res.*, **98**, 20737-20759.
- Beck, C., J. Grieser and B. Rudolf, 2005: A new monthly precipitation climatology for the global land areas for the period 1951 to 2000. *Klimastatusbericht 2004*, DWD, ISSN 1437-7691, ISSN 1616-5063 (Internet), ISBN 3-88148-402-7, 181-190.
- Beyer, H.G., C. Costanzo, and D. Heinemann, 1996: Modifications of the Heliosat procedure for irradiance estimates from satellite images. *Solar Energy*, **56**, 207-212.
- Brohan, P. *et al.*, 2006: Uncertainty estimates in regional and global observed temperature changes. *J. Geophys. Res.*, **111**, D12106.
- Brunet M., P. D. Jones, J. Sigró, O. Saladié, E. Aguilar, A. Moberg, P. M. Della-Marta, D. Lister, A. Walther, and D. López, 2007a: Temporal and spatial temperature variability and change over Spain during 1850-2005. *J. Geophys. Res.*, **112**, D12117, doi:10.1029/2006JD008249.
- Brunet M., J. Sigró, P.D. Jones, O. Saladié, E. Aguilar, A. Moberg, D. Lister and A. Walther, 2007b: Long-term changes in extreme temperatures and precipitation in Spain, *Contributions to Science*, **3**, Barcelona, 331-342.



- Brunet, M., O. Saladié, P. Jones, J. Sigró, E. Aguilar, A. Moberg, D. Lister, A. Walther, and C. Almarza, 2008: A case-study/Guidance on the development of long-term daily adjusted temperature datasets. WMO-TD-1425 - WCDMP-66, Geneva: 43 pp.
- Brunet, M. and F. G. Kuglitsch (Eds.), 2008: Proceedings of the International Workshop on Rescue and Digitization of Climate Records in the Mediterranean Basin, WCDMP No. 67/WMO-TD No. 1432, XXII + 219 pp.
- Butler, D., 2007: The planetary panopticon. *Nature*, **450**, 778-781.
- Deneke, H.M., A.J. Feijt, and R.A. Roebeling, 2008: Estimating surface solar irradiance from METEOSAT SEVIRI-derived cloud properties. *Remote Sens. Environ.*, **112**, 3131-3141.
- Drews, A., H.G. Beyer and U. Rindelhardt, 2008: Quality of performance assessment of PV plants based on irradiance maps. *Solar Energy*, **82**, 1067-1075, doi:10.1016/j.solener.2008.04.009
- Dürr, B. and A. Zelenka, 2008: Deriving surface global irradiance over the Alpine region from METEOSAT Second Generation data by supplementing the HELIOSAT method. *International Journal of Remote Sensing* (in press).
- Dürr, B., A. Zelenka, R. Müller and R. Philipona, 2008: Verification of CM-SAF and MeteoSwiss satellite based retrievals of surface shortwave irradiance over the Alpine region. *International Journal of Remote Sensing* (accepted).
- Durand, Y., M. Laternser, G. Giraud, P. Etchevers, B. Lesaffre, and L. Mérindol, 2008: Reanalysis of 44 years of climate in French Alps (1958-2002); methodology, model validation, climatology and trends for air temperature and precipitation. *J. of App. Met. and Clim.* Doi:10.1175/2008JAMC1808.1 (submitted).
- Durre, I., R.S. Vose, and D.B. Wuertz, 2006: Overview of the integrated global radiosonde archive. *J. Climate.*, **19**, 53-68.
- EC, 2008: Global Monitoring for Environment and Security (GMES): we care for a safer planet. Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. COM(2008) 748 final.
- EEA-JRC-WHO, 2008: Impacts of Europe's changing climate – 2008 indicator-based assessment. EEA Report No 4/2008, JRC Reference Report No JRC47756, DOI 10.2800/48117.
- Fortelius, C., A. Andrae, and M. Forsblomi, 2002: The BALTEX regional reanalysis project. *Boreal Env. Res.*, **7**, 193-201.
- Frei, C. and C. Schär, 1998: A precipitation climatology of the Alps from high-resolution rain-gauge observations. *Int. J. Climatol.*, **18**, 873-900.
- GCOS, 2003: GCOS Second Report on the Adequacy of the Global Climate Observing System in Support of the UNFCCC. GCOS-92, WMO/TD No. 1143, 75pp.
- GCOS, 2004: GCOS Implementation Plan for Global Observing System for Climate in support of UNFCCC. GCOS-92, WMO/TD No. 1219, 136 pp.
- GEO, 2007: GEO Group on Earth Observations Work Plan for 2007-2009. March 2007. 37pp., [www.earthobservations.org/documents/wpo709\\_v4.pdf](http://www.earthobservations.org/documents/wpo709_v4.pdf)
- Grassl, H., V. Jost, R. Kumar, J. Schulz, P. Bauer, P. Schluessel, 2000: The Hamburg Ocean-Atmosphere Parameters and Fluxes from Satellite Data (HOAPS): A Climatological Atlas of Satellite-Derived Air-Sea-Interaction Parameters over the Oceans., Report No. 312, ISSN 0937-1060, Max Planck Institute for Meteorology, Hamburg.
- Gustafsson, N., L. Berre, S. Hörnquist, X.Y. Xuang, M. Lindskog, B. Navascues, K.S. Mogensen, and S. Thorsteinsson, 2000: Three-dimensional variational data assimilation for a limited area model, Part I: General formulation and the background error constraint. *Tellus*, **53A**, 425-446.

- Habets, F., A. Boone, J.L. Champeaux, P. Etchevers, L. Franchisteguy, E. Leblois, E. Ledoux, P. Le Moigne, E. Martin, S. Morel, J. Noilhan, P. Quintana Segui, F. Rousset-Regimbeau and P. Viennot, 2008: The SAFRAN-ISBA-MODCOU hydrometeorological model applied over France. *J. Geophys. Res.*, **113**, D06113, doi:10.1029/2007JD008548.
- Haimberger, L., 2005: Homogenization of radiosonde temperature time series using ERA-40 analysis feedback information. ERA-40 Project Report Series 23, ECMWF, Reading, U.K., 68 pp.
- Haimberger, L., C. Tavalato, and S. Sperka, 2008: Toward elimination of the warm bias in historic radiosonde temperature records - Some new results from a comprehensive intercomparison of upper-air data. *J. Climate*, **21**, 4587–4606, DOI: 10.1175/2008JCLI1929.1.
- Hammer, A., D. Heinemann, C. Hoyer, R. Kuhlemann, E. Lorenz, R.W. Müller, H.G. Beyer, 2003: Solar energy assessment using remote sensing technologies. *Remote Sensing of Environment*, **86**, 423-432.
- Haylock, M.R., N. Hofstra, A.M.G. Klein Tank, E.J. Klok, P.D. Jones, M. New, 2008: A European daily high-resolution gridded dataset of surface temperature and precipitation. *J. Geophys. Res.*, **113**, doi:10.1029/2008JD10201.
- Hofstra, N. M. Haylock, M. New, P. Jones, and C. Frei, 2008: Comparison of six methods for the interpolation of daily, European climate data. *J. Geophys. Res.*, **113**, doi:10.1029/2008JD010100.
- Jansson, A., C. Persson, and G. Strandberg, 2007: 2D meso-scale re-analysis of precipitation, temperature and wind over Europe – ERAMESAN. Time period 1980-2004. SMHI, RMK Report No 112.
- Jenkins, G.J., M.C. Perry, and M.J.O. Prior, 2007: The climate of the United Kingdom and recent trends. Met Office Hadley Centre, Exeter, UK.
- Kabat, P. and P. Velinga, 2005: Climate proofing the Netherlands. *Nature*, **438**, 283-284.
- Klein Tank, A.M.G., T.C. Peterson, D.A. Quadir, S. Dorji, X. Zou, H. Tang, K. Santhosh, U.R. Joshi, A.K. Jaswal, R.K. Kolli, A. Sikder, N.R. Deshpande, J.V. Revadekar, K. Yeleuova, S. Vandasheva, M. Faleyeva, P. Gomboluudev, K.P. Budhathoki, A. Hussain, M. Afzaal, L. Chandrapala, H. Anvar, D. Amanmurad, V.S. Asanova, P.D. Jones, M.G. New, T. Spektorman, 2006: Changes in daily temperature and precipitation extremes in central and south Asia. *J. Geophys. Res.*, **111** (D16105), doi:10.1029/2005JD006316.
- Li, J., W.W. Wolf, W.P. Menzel, W. Zhang, H.-L. Huang, and T.H. Achtor, 2000: Global soundings of the atmosphere from ATOVS measurements: The algorithm and validation. *J. Appl. Meteor.*, **39**, 1248-1268.
- Lindskog, M., N. Gustafsson, B. Navascués, K.S. Mogensen, X.-Y. Huang, X. Yang, U. Andrae, L. Berre, S. Thorsteinsson, and J. Rantakokko, 2001: Three-dimensional variational data assimilation for a limited area model. Part II: Observation handling and assimilation experiments. *Tellus*, **53A**, 447-468.
- Mesinger, F., *et al.*, 2006: North American Regional Reanalysis. *BAMS*, DOI:10.1175/BAMS-87-3-343.
- Mitchell, T.D., and P.D. Jones, 2005: An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *Int. J. Climatol.*, **25**, 693–712.
- Moberg, A., P. D. Jones, D. Lister, A. Walther, M. Brunet, J. Jacobeit, L. V. Alexander, P. M. Della-Marta, J. Luterbacher, P. Yiou, D. Chen, A. M. G. Klein Tank, O. Saladié, J. Sigró, E. Aguilar, H. Alexandersson, C. Almarza, I. Auer, M. Barriendos, M. Begert, H. Bergström, R. Böhm, C. J. Butler, J. Caesar, A. Drebs, D. Founda, F. W. Gerstengarbe, G. Micela, M. Maugeri, H. Österle, K. Pandzic, M. Petrakis, L. Srnec, R. Tolasz, H. Tuomenvirta, P. C. Werner, H. Linderholm, A. Philipp, H. Wanner, and E. Xoplaki, 2006: Indices for daily temperature and precipitation extremes in Europe analysed for the period 1901-2000. *J. Geophys. Res.*, **111** (D22106), doi:10.1029/2006JD007103.

- Navascuès, B., 1997: Analysis of 2 meter Temperature and Relative Humidity. HIRLAM Tech. Report No. 28, S-601 76, SMHI, Norrköping, SWEDEN.
- Parker, D.E., 2004: Large scale warming is not urban. *Nature*, **432**, 290.
- Peterson, T. C., *et al.*, 2001: Report on the activities of the Working Group on Climate Change Detection and Related Rapporteurs 1998-2001. World Meteorological Organisation Rep. WCDMP-47, WMO-TD 1071, Geneva, Switzerland, 143 pp.
- Peterson T. C., X. Zhang, M. Brunet, and J.L. Vázquez Aguirre, 2007: Changes in North American extremes derived from daily weather data. *J. Geophys. Res.*, doi:10.1029/2007JD009453.
- Peterson, T.C., and M.J. Manton, 2008: Monitoring changes in climate extremes - A tale of international collaboration. *Bull. Amer. Meteorol. Soc.*, DOI:10.1175/2008BAMS2501.1.
- Rayner, N.A., *et al.*, 2006: Improved analyses of changes and uncertainties in marine temperature measured *in situ* since the mid-nineteenth century: the HadSST2 dataset. *J. Climate*, **19**, 446-469.
- Rodríguez E., B. Navascuès, J.J. Ayuoso, and S. Järvenoja, 2003: Analysis of surface variables and parameterization of surface processes in HIRLAM. Part 1: Approach and verification by parallel runs. HIRLAM Tech. Report No. 58, S-601 76, SMHI, Norrköping, SWEDEN.
- Roebeling, R.A., A.J. Feijt, and P. Stammes, 2006: Cloud property retrievals for climate monitoring: implications of differences between SEVIRI on METEOSAT-8 and AVHRR on NOAA-17. *J. Geophys. Res.*, **111**, D20210, doi:10.1029/2005JD006990.
- Roebeling, R.A. and I. Holleman, 2008: Validation of rain rate estimates from SEVIRI using weather radar observations, Proceedings EUMETSAT Meteorological Satellite Conference, 8 – 12 September 2008, Darmstadt, Germany.
- Rudolf, B., and U. Schneider, 2005: Calculation of Gridded Precipitation Data for the Global Land-Surface using in situ Gauge Observations. Proceedings of the 2nd Workshop of the International Precipitation Working Group IPWG, Monterey October 2004, EUMETSAT, ISBN 92-9110-070-6, ISSN 1727-432X, 231-247.
- Schulz, J, P. Albert, H.-D. Behr, D. Caprion, H. Deneke, S. Dewitte, B. Duerr, P. Fuchs, A. Gratzki, P. Hechler, R. Hollmann, S. Johnston, K.-G. Karlsson, T. Manninen, R. Müller, M. Reuter, A. Riihelä, R. Roebeling, N. Selbach, A. Tetzlaff, W. Thomas, M. Werscheck, E. Wolters, and A. Zelenka, 2008: Operational climate monitoring from space: the EUMETSAT satellite application facility on climate monitoring (CM-SAF), *Atmos. Chem. Phys. Discuss.*, **8**, 8517–8563.
- Simmons, A.J., P.D. Jones, V. da Costa Bechtold, A.C.M. Beljaars, P.W. Kållberg, S. Saarinen, S.M. Uppala, P. Viterbo and N. Wedi, 2004: Comparison of trends and low-frequency variability in CRU, ERA-40 and NCEP/NCAR analyses of surface air temperature. *J. Geophys. Res.*, **109**, D24115, doi:10.1029/2004JD006306.
- Thorne, P.W., *et al.*, 2005a: Revisiting radiosonde upper-air temperatures from 1958 to 2002. *J. Geophys. Res.*, **110**, D18105, doi:10.1029/2004JD00575.
- Thorne, P.W., Parker, D.E., Christy, J.R. and Mears, C.A., 2005b: Uncertainties in climate trends: lessons from upper-air temperature records. *Bull. Amer. Meteorol. Soc.*, **86**, 1437-1442.
- Titchner, H., M.P. McCarthy, P.W. Thorne, S.F.B. Tett, L. Haimberger, and D.E. Parker, 2008: Critically reassessing tropospheric temperature trends from radiosondes using realistic validation experiments. *J. Climate* (in press).
- Trentmann, J. and R.W. Mueller, *et al.*, 2008: Evaluation of ERA-interim downwelling surface radiation fluxes with BSRN surface observations, DWD (manuscript in preparation).

- Undén, P., L. Rontu, H. Järvinen, P. Lynch, J. Calvo, G. Cats, J. Cuxart, K. Eerola, C. Fortelius, J. Antonio Garcia-Moya, C. Jones, G. Lenderlink, A. McDonald, R. McGrath, B. Navascues, N. Woetman Nielsen, V. Ødegaard, E. Rodriguez, M. Rummukainen, R. Rööm, K. Sattler, B. Hansen Sass, H. Savijärvi, B. Wichers Schreur, R. Sigg, H. The, A. Tijm, 2002: HIRLAM-5 Scientific Documentation, HIRLAM-5 Project. c/o Per Undén SMHI, S-601 76 Norrköping, SWEDEN.
- Uppala, S.M., P. Källberg, A.J. Simmons, U. Andrae, V. Da Costa Bechtold, M. Fiorino, K. Gibson, J. Haseler, A. Hernandez., G.A. Kelly, K. Li, K. Onogi, S. Saarinen, N. Sokka, R.P. Allan, E. Andersson, K. Arpe, M. Balmaseda, A.C.M. Beljaars, L. Van De Berg, J. Bidlot, N. Bormann, S. Caires, F. Chevallier, A. Dethof, M. Dragosavac, M. Fisher, M. Fuentes, S. Hagemann, E. Hólm, B.J. Hoskins, L. Isaksen, P.A.E.M. Janssen, R. Jenne, A.P. McNally, J.-F. Mahfouf, J.-J. Morcrette, N.A. Rayner, R.W. Saunders, P. Simon, A. Sterl, K.E. Trenberth, A. Untch, D. Vasiljevic, P. Viterbo, and J. Woollen, 2005: The ERA-40 reanalysis. *Quart. J. Roy. Meteor. Soc.*, **131**, 2961-3012.
- Wang, L. and J.J. Qu, 2007: NMDI: A normalized multi-band drought index for monitoring soil and vegetation moisture with satellite remote sensing, *Geophys. Res. Letters*, **34**, L20405, doi:10.1029/2007GL031021.
- Wang K. and S. Liang, 2008: An improved method for estimating global evapotranspiration based on satellite determination of surface net radiation, vegetation index temperature and soil moisture. *Journal of Hydrometeorology*, **9**, 712-727.
- Wijngaard, J.B., A.M.G. Klein Tank, and G.P. Können, 2003: Homogeneity of 20th century European daily temperature and precipitation series. *Int. J. Climatol*, **23**, 679-692.
- Willett, K.M., N.P. Gillett, P.D. Jones and P.W. Thorne, 2007: Attribution of observed surface humidity changes to human influence. *Nature*, **449**, 710-713.
- Willett, K.M., P.D. Jones, N.P. Gillett and P.W. Thorne, 2008: Recent changes in surface humidity: Development of the HadCRUH dataset. *J. Climate*, **21**, 5364-5383.
- WMO, 2005: The WCRP Strategic Framework 2005-2015 Coordinated Observation and Prediction of the Earth System (COPES). WCRP-123, WMO/TD-No. 1291, August 2005, 59 pp.
- Zhang, X., E. Aguilar, S. Sensoy, H. Melknyan, U. Taghiyeva, N. Ahmed, N. Kutaladze, F. Rahimzadeh, A. Taghipour, T. H. Hantosh, P. Albert, M. Semawi, M.K. Ali, M. Halal, S. Al-Shabibi, Z. Al-Oulan, T. Zatari, I. Al Dean Khalil, R. Sagir, M. Demircan, M. Eken, M. Adiguzel, L. Alexander, T.C. Peterson and T. Wallis, 2005: Trends in Middle East climate extremes indices during 1930-2003. *J. Geophys. Res.*, **110**, D22104, doi:10.1029/2005JD006181.

## Appendix A: Acronym list

ACRE: Atmospheric Circulation Reconstructions over the Earth  
AIREP: Air Report (aviation)  
ALADIN: Limited-area NWP model  
ALP-IMP: Multi-centennial climate variability in the Alps based on Instrumental data, Model simulations and Proxy data  
AMDAR: Aircraft Meteorological Data Relay  
AOPC: Atmospheric Observation Panel for Climate (part of GCOS)  
AROME: High resolution model of Météo France  
ARPEGE-IFS: atmospheric model developed jointly by Météo France and ECMWF  
ATOVs: Advanced TIROS Operational Vertical Sounder  
BALTEX: Baltic Sea Experiment  
BSRN: Baseline Surface Radiation Network  
CarboEurope: Assessment of the European Terrestrial Carbon Balance  
CCI: Commission for Climatology (of WMO)  
CECILIA: Central and Eastern Europe Climate Change Impact and Vulnerability Assessment (EU-project)  
CIB: Climate Information Bulletin (part of EURO4M)  
CIRCE: Climate change and impact research: the Mediterranean environment (EU-project)  
CIRCLE: Climate Impact Research Coordination within a Larger Europe (EU ERA-Net activity)  
CIRES: Cooperative Institute for Research in Environmental Sciences  
CLARIS-LPB: A Europe-South America Network for Climate Change Assessment and Impact Studies  
CLIVAR: International research programme on Climate Variability and Predictability  
CLIWOC: Climate of the world's oceans (EU-project)  
CLT: Climate Liaison Team (part of EURO4M)  
CM-SAF: Satellite Application Facility on Climate Monitoring  
COPES: Coordinated Observation and Prediction of the Earth System (WCRP strategy 2005-2015)  
COST: European Co-operation in the field of Scientific and Technical Research  
CPP: Cloud Physical Properties  
CRUTEM3: Climatic Research Unit's Land Air Temperature database  
DARE: Data Rescue (WMO initiative)  
DWD: Deutscher Wetterdienst (German Weather Service)  
DYNAMITE: Understanding the Dynamics of the Coupled Climate System (EU-project)  
E-Obs: ENSEMBLES gridded daily dataset for Europe  
ECA&D: European Climate Assessment & Dataset  
ECMWF: European Centre for Medium-Range Weather Forecasts  
ECOCHANGE: Biodiversity and ecosystem changes in Europe (FP-project)  
ECVs: Essential Climate Variables (defined by GCOS)  
EEA: European Environment Agency  
EIONET: European Environment Information and Observation NETWORK

ELDAS: Development of a European Land Data Assimilation System to predict Floods and Droughts (EU-project)  
 EMI: European Meteorological Infrastructure  
 EMSLP: EMULATE Mean Sea Level Pressure dataset  
 EMULATE: European and North Atlantic daily to MULTidecadal climATE variability (EU project)  
 ENSEMBLES: ENSEMBLE-based Predictions of Climate Changes and their Impacts (EU-project)  
 EQUAL II: A laboratory for new ideas to the European Employment Strategy and the Social inclusion process (EU initiative)  
 ERA-Interim: ECMWF reanalysis  
 ERA-40: ECMWF reanalysis for the period from September 1957 to August 2002  
 ERA-75: Proposed next generation ECMWF reanalysis  
 ERA-CLIM: Development of a European Reanalysis Capability for Global Climate Monitoring (FP7-proposal)  
 ERAMESAN: Meso-scale reanalysis by SMHI  
 ERA-Net: FP-scheme for the coordination and cooperation of national and regional programmes  
 ERBE: Earth Radiation Budget Experiment  
 ESA: European Space Agency  
 ESPON: European Spatial Planning Observation Network  
 ESRL: Earth System Research Laboratory (USA)  
 ETCCDI: Expert Team on Climate Change Detection and Indices  
 EUMETNET-ECSN: European Climate Support Network of the Network of European Meteorological Services  
 EUMETSAT: European Meteorological Satellite Organisation  
 EURO-LIMPACS: Integrated project to evaluate impacts of global change on European freshwater ecosystems (EU-project)  
 EURO4M: European Reanalysis and Observations for Monitoring (this project)  
 EuroClim: European Climate Change Monitoring and Prediction System (EU-project)  
 EuroCryoClim: European Climate Change Monitoring and Prediction System  
 EUROGRID: European Gridding (EUMETNET-ECSN project)  
 EuroGLOBEC: European global ocean ecosystem dynamics  
 EURRA: European Regional Reanalysis  
 GAIA-network: Network for female earth scientists in the Netherlands  
 GAR: Greater Alpine Region  
 GAW: Global Atmospheric Watch  
 GCOS: Global Climate Observing System  
 GEMS: Global and regional Earth-system (Atmosphere) Monitoring using Satellite and *in situ* data  
 GEO: Group on Earth Observations  
 GEOLAND2: Integrated GMES Project On Land Cover and Vegetation (EU-project)  
 GEOSS: Global Earth Observations System of Systems  
 GERB/CERES: Geostationary Earth Radiation Budget/Clouds and the Earth's Radiant Energy System  
 GEWEX: Global Energy and Water Cycle Experiment  
 GIS: Geographical Information System  
 GMES: Global Monitoring for Environment and Security  
 GMR: Greater Mediterranean Region

GPCC: Global Precipitation Climatology Centre  
 GPCP: Global Precipitation Climatology Project  
 GSN: GCOS Surface Network  
 GTS: Global Telecommunications System  
 GUAN: GCOS Upper Air Network  
 HadAT: Hadley Centre gridded radiosonde temperature dataset  
 HadCRUH: Hadley Centre and Climatic Research Unit's monthly mean surface humidity dataset  
 HadCRUT<sub>3</sub>: Hadley Centre and Climatic Research Unit's Air and Marine Temperature Anomalies Version 3 dataset  
 HadISST<sub>2</sub>: Hadley Centre Sea Ice and Sea Surface Temperature dataset  
 HadSLP: Hadley Centre mean Sea Level Pressure dataset  
 HadSST<sub>2</sub>: Hadley Centre Sea Surface Temperature dataset  
 HIRLAM: High-Resolution Limited-Area Model  
 HOAPS: Hamburgs Ocean Atmosphere Parameters from Satellite dataset  
 HOLSMEER: Late holocene shallow marine environments of Europe (FP-project)  
 HOME: Advances in Homogenisation Methods of Climate Series: An Integrated Approach (COST-Action)  
 HRT-GAR: High resolution Temperature Climatology for the Greater Alpine Region  
 H-SAF: Satellite Application Facility on Support to Operational Hydrology and Water Management  
 IAPP: International ATOVS Processing Package  
 ICOADS: International Comprehensive Ocean-Atmosphere Dataset  
 IGRA: Integrated Global Radiosonde Archive ([www.ncdc.noaa.gov/oa/cab/igra/index.php](http://www.ncdc.noaa.gov/oa/cab/igra/index.php))  
 IMPROVE: Improved Understanding of past climatic variability from early daily European instrumental sources (EU-project)  
 INSPIMET: EUMETNET group that oversees the user requirements for the implementation rules of INSPIRE  
 INSPIRE: INfrastructure for SPatial InfoRmation in Europe  
 INTAS: International Association for the Promotion of Co-operation with Scientists from the New Independent States of the Former Soviet Union  
 IPCC: Intergovernmental Panel on Climate Change  
 IPCC-AR<sub>4</sub>: Fourth Assessment Report of the IPCC  
 IPCC-TAR: Third Assessment Report of the IPCC  
 IPY-CARE: Climate of the Arctic and its Role for Europe (CARE) - a European component of the International Polar Year (EU-project)  
 ISCCP: International Satellite Cloud Climatology Data  
 ISH: Integrated Surface Hourly dataset  
 ISOWG: GMES In Situ Observation Working Group  
 ISPD: International Surface Pressure Data Bank  
 JCOMM: Joint WMO/IOC Commission for Oceanography and Marine Meteorology  
 JRC: Joint Research Centre  
 KNMI: Royal Netherlands Meteorological Institute  
 Land-SAF: Land Surface Analysis Satellite Applications Facility  
 LIFE AIRFORALL: Air Pollution Forecasting, Alert and Monitoring System on Short Time Scale, at local and regional scale in unfavourable meteorological and topographic conditions  
 LIFE ASSURE: Assessment System for Urban Environment

MACC: Monitoring Atmospheric Composition and Climate  
 MEDARE: MEditerranean climate DAta REscue Initiative under the auspice of WMO  
 MedCLIVAR: Mediterranean CLIVAR  
 MESAN: Meso-scale analysis system by SMHI  
 METAFOR: Common Metadata for Climate Modelling Digital Repositories (EU-project)  
 METEOSAT: Meteorological Satellite of ESA and EUMETSAT  
 MF: Météo France  
 MICE: Modelling the Impact of Climate Extremes (EU-project)  
 MO: Met Office  
 MOCCA: MOonitoring Climate variability and Change for an improved environmental and risk management in the Alpine space  
 MOODLE: course management system (Open Source)  
 MS: Meteo Swiss  
 MSLP: Mean Sea Level Pressure  
 MSG: METEOSAT Second Generation  
 MSU: Microwave Sounding Unit  
 MVIRI (SEVIRI): satellite instrument on-board of MSG  
 MyOCEAN: GMES service for ocean monitoring and forecasting  
 NAE: North Atlantic and European (NWP configuration)  
 NARR: North American Regional Reanalysis  
 NCAR: National Center for Atmospheric Research (Boulder, Co, USA)  
 NCDC: National Climatic Data Center (Asheville, NC, USA)  
 NCEP: National Centers for Environmental Prediction (USA)  
 NERC: Natural Environmental Research Council (UK)  
 NetFAM: Nordic Network on Fine-scale Atmospheric Modelling  
 NitroEurope: Contribution of Nitrogen to the net greenhouse gas budgets of Europe  
 NMA: National Meteorological Administration  
 NMHS: National Meteorological and Hydrological Service  
 NOAA: National Oceanic & Atmospheric Administration (USA)  
 NORDKLIM: Nordic co-operation within climate activities  
 NWP: Numerical Weather Prediction  
 OOPC: Oceanic Observation Panel for Climate (part of GCOS)  
 OPAG: Open Programme Area Group (WMO/CCL body)  
 QC: Quality Control  
 QCCCE: Queensland Climate Change Centre of Excellence (Australia)  
 PDSI: Palmer Drought Severity Index  
 PET: Potential Evapotranspiration  
 PROMOTE: PROtocol MOniToring for the GMES Service Element: Atmosphere  
 PRUDENCE: Prediction of Regional scenarios and Uncertainties for Defining EuropeaN Climate change risks and Effects (FP-project)  
 SAF: Satellite Application Facility (EUMETSAT)  
 SAFRAN: Meteorological analysis system  
 SBSTA: Subsidiary Body for Scientific and Technological Advice  
 SCCONE: Snow Cover Changes over Northern Eurasia during the last century (INTAS-project)  
 SCIAMACHY: SCanning Imaging Absorption SpectroMeter for Atmospheric CHartographY



SDI: Spatial Data Infrastructure  
SEIS: Shared Environmental Information System  
SEVIRI: satellite instrument on-board of MSG  
SIM: Hydrometeorological model  
SMHI: Swedish Meteorological and Hydrological Institute  
SST: Sea Surface Temperature  
STARDEX: Statistical and Regional dynamical Downscaling of Extremes for European regions (EU project)  
SYNOP: Synoptic observation  
UEA: University of East Anglia (Climatic Research Unit)  
UKCIP08: UK 21st Century Climate Scenarios  
UNESCO-Bilko: United Nations Educational, Scientific and Cultural Organization project for training in coastal and marine remote sensing  
UNFCCC: United Nations Framework Convention on Climate Change  
URV: University Rovira i Virgili  
WASA: Waves and Storms in the North Atlantic (FP-project)  
WCDMP: World Climate Data Monitoring Programme  
WCRP: World Climate Research Programme  
WEELS: Wind Erosion on European Light Soils (FP-project)  
WHO: World Health Organization  
WIS: WMO Information System  
WMO: World Meteorological Organization  
WWW: World Weather Watch  
3D-Var: Three-dimensional Variational (analysis or assimilation)  
4D-Var: Four-dimensional Variational (analysis or assimilation)

## Appendix B: Support letters

01/12 '08 16:26 FAX 33367151

EEA EAS

001/001

European Environment Agency



Dr Albert Klein Tank  
KNMI  
3730 AE De Bilt  
The Netherlands

28 November 2008  
AJO

*Dear Dr Klein Tank,*

**'Letter of support' to the proposed FP7 project EURO4M**

We have been informed that a consortium of partners from the scientific community will submit a proposal 'European Reanalysis and Observations for Monitoring (EURO4M)' to the EU Seventh Framework Programme Cooperation Theme 'Space', aiming at significant improvements of climate change atmospheric time series and monitoring services for Europe.

The European Environment Agency (EEA) as a potential user of EURO4M products is very pleased to offer support in order to guide the project towards its very important objectives. The integrated usage of physical models describing atmospheric motion together with a variety of different sources of observational data is expected to deliver gridded data sets highly relevant to environmental assessments, in particular for regional climate impact assessments, and the development of cost-effective adaptation strategies. The resulting detailed information on past climate change will allow to better identify vulnerable sectors and regions with a need for adaptation.

Especially for regional and local assessments, Europe currently still lacks data of appropriate spatial resolution and quality. Adding on to the existing GMES core service components, EURO4M's efforts to provide comprehensive pan-European climate datasets will help to address these critical gaps in data coverage.

*With best regards*

A handwritten signature in black ink, appearing to read 'J. McGlade'.

Prof. Jacqueline McGlade  
Executive Director

Kongens Nytorv 6  
1050 Copenhagen K  
Denmark

Tel.: +45 33 36 71 00  
Fax: +45 33 36 71 99

E-mail: [eea@eea.europa.eu](mailto:eea@eea.europa.eu)  
Web: [www.eea.europa.eu](http://www.eea.europa.eu)



2<sup>nd</sup> December 2008

Dr Albert Klein Tank  
KNMI  
3730 AE De Bilt  
The Netherlands

**EUMETNET's "Letter of support" to the proposed FP7 project EURO4M**

Dear Dr Klein Tank

EUMETNET, the Network of 24 European Meteorological Services, is aware of the EURO4M proposal which will be submitted to the EU Seventh Framework Programme Cooperation Theme "Space" aiming to develop a European capacity to provide and deliver the best possible climate change time series and monitoring services for European users.

This is a very important project that will provide data products and climate change services to a diverse user community, from policy-makers to European citizens and will help them to adapt to climate variability and climate change at the regional, sub-regional and national scales.

EUMETNET and its Members are active in the fields of climate monitoring and climate change. EUMETNET's European Climate Support Network (ECSN) Programme aims at providing sustainable high quality and climate reference services, tailored to the users needs. EUMETNET supports this proposal, because EURO4M will integrate with EUMETNET-ECSN on the production of gridded climate data sets based on dense networks of station observations, particularly in the next EUROGRID proposal which is in preparation by ECSN members for the EUMETNET Council (as a follow-up to S-EUROGRID).

EURO4M consortium participants are strategically placed to liaise with a wide range of international and national bodies. All European NMHSs are linked to EURO4M via its close association with ECMWF, EUMETNET, EUMETSAT, and their joint European Meteorological Infrastructure (EMI).

Yours sincerely,

Steve Noyes  
Executive Director  
EUMETNET

EUMETNET Coordination Office  
c/o The Royal Meteorological Institute of Belgium, Ringlaan 3 Avenue Circulaire, 1180 Brussels, Belgium  
Phone : +32 (0)2 373 05 18 Fax : +32 (0)2 890 98 58